THE SUMMER MEETING IN HANOVER

The forty-sixth Summer Meeting of the Society and the twenty-second Colloquium were held at Dartmouth College, Hanover, New Hampshire, from Tuesday to Thursday, September 10–12, 1940. The Mathematical Association of America met on Monday morning and afternoon and the Institute of Mathematical Statistics on Tuesday morning and afternoon and Wednesday and Thursday mornings. About four hundred fifty persons attended the meetings, among whom were the following two hundred ninety-eight members of the Society:

C. R. Adams, R. B. Adams, R. P. Agnew, Leon Alaoglu, Warren Ambrose, R. C. Archibald, Herbert E. Arnold, H. T. R. Aude, R. P. Bailey, J. L. Barnes, R. C. F. Bartels, Felix Bernstein, R. A. Beth, E. E. Betz, Harry Birchenough, Garrett Birkhoff, G. D. Birkhoff, R. P. Boas, Salomon Bochner, C. B. Boyer, A. T. Brauer, R. W. Brink, B. H. Brown, B. L. Brown, R. E. Bruce, G. S. Bruton, J. A. Bullard, C. T. Bumer, R. S. Burington, F. J. H. Burkett, J. H. Bushey, Jewell H. Bushey, W. D. Cairns, B. H. Camp, M. E. Carlen, Leonard Carlitz, W. B. Carver, J. A. Clarkson, R. F. Clippinger, L. W. Cohen, Nancy Cole, Esther Comegys, G. M. Conwell, T. F. Cope, A. H. Copeland, L. P. Copeland, Richard Courant, A. P. Cowgill, A. T. Craig, A. R. Crathorne, H. B. Curry, D. R. Curtiss, J. H. Curtiss, Tobias Dantzig, D. R. Davis, F. F. Decker, L. S. Dederick, M. R. Demers, A. H. Diamond, C. E. Dimick, C. H. Dix, W. J. Dixon, H. A. DoBell, J. L. Doob, T. L. Downs, Arnold Dresden, John Dyer-Bennet, Samuel Eilenberg, Churchill Eisenhart, L. P. Eisenhart, M. L. Elveback, Benjamin Epstein, Paul Erdös, G. C. Evans, G. W. Evans, W. K. Feller, W. B. Fite, H. T. Fleddermann, M. M. Flood, L. R. Ford, G. E. Forsythe, Tomlinson Fort, R. M. Foster, A. H. Fox, J. S. Frame, Philip Franklin, K. O. Friedrichs, Orrin Frink, T. C. Fry, R. E. Fullerton, M. G. Gaba, H. L. Garabedian, H. J. Gay, H. P. Geiringer, B. P. Gill, R. E. Gilman, Michael Goldberg, Michael Golomb, C. H. Graves, L. J. Green, E. H. Hadlock, D. W. Hall, Marshall Hall, P. R. Halmos, Israel Halperin, J. A. Hamilton, D. C. Harkin, O. G. Harrold, O. C. Hazlett, E. R. Hedrick, M. H. Heins, Robert Henderson, J. G. Herriot, E. H. C. Hildebrandt, T. H. Hildebrandt, Einar Hille, M. P. Hollcroft, T. R. Hollcroft, G. M. Hopper, Harold Hotelling, S. E. Hotelling, E. M. Hull, E. V. Huntington, W. A. Hurwitz, M. H. Ingraham, Dunham Jackson, S. B. Jackson, Fritz John, R. A. Johnson, F. E. Johnston, A. W. Jones, B. W. Jones, H. A. Jordan, Mark Kac, M. L. Kales, E. R. van Kampen, Wilfred Kaplan, E. S. Kennedy, L. S. Kennison, S. C. Kleene, J. R. Kline, J. C. Knipp, E. R. Kolchin, W. D. Lambert, O. E. Lancaster, A. E. Landry, R. E. Langer, R. A. Leibler, B. A. Lengyel, C. A. Lester, Norman Levinson, D. C. Lewis, M. I. Logsdon, W. R. Longley, L. H. Loomis, C. I. Lubin, N. H. McCoy, E. J. McShane, L. A. MacColl, C. C. MacDuffee, Saunders MacLane, H. M. MacNeille, W. G. Madow, Dorothy Manning, L. C. Mathewson, Walther Mayer, W. E. Milne, Richard von Mises, E. B. Mode, Deane Montgomery, T. W. Moore, F. M. Morgan, W. K. Morrill, Max Morris, Marston Morse, E. J. Moulton, G. W. Mullins, C. W. Munshower, O. E. Neugebauer, John von Neumann, C. V. Newsom, A. V. Newton, Jerzy Neyman, C. O. Oakley, R. E. O'Connor, P. S. Olmstead, J. M. H. Olmsted, Isaac Opatowski, Oystein Ore, F. W. Owens, H. B. Owens, Gordon Pall, E. W. Paxson, F. W. Perkins, O. J. Peterson, C. G. Phipps, Everett Pitcher, Hillel

Poritsky, G. B. Price, E. J. Purcell, H. A. Rademacher, E. D. Rainville, J. F. Randolph, M. S. Rees, R. W. Rempfer, Harris Rice, R. G. D. Richardson, D. E. Richmond, R. F. Rinehart, J. F. Ritt, H. E. Robbins, Robin Robinson, S. L. Robinson, Benjamin Rosenbaum, Joseph Rosenbaum, R. A. Rosenbaum, Arthur Rosenthal, J. B. Rosser, H. G. Russell, N. E. Rutt, S. T. Sanders, Peter Scherk, E. R. Schneckenburger, I. J. Schoenberg, G. E. Schweigert, C. H. W. Sedgewick, R. W. Sedgewick, W. E. Sewell, I. M. Sheffer, L. W. Sheridan, L. L. Silverman, R. G. Simond, M. E. Sinclair, Abraham Sinkov, H. L. Slobin, M. M. Slotnick, M. F. Smiley, C. E. Smith, P. A. Smith, T. L. Smith, W. M. Smith, Virgil Snyder, M. R. Solt, Joseph Spear, D. C. Spencer, V. E. Spencer, Abraham Spitzbart, G. R. Stibitz, C. N. Stokes, M. H. Stone, E. C. Strayhorn, R. E. Street, D. J. Struik, Otto Szász, Gabor Szegö, J. D. Tamarkin, J. M. Thomas, E. W. Titt, M. M. Torrey, J. I. Tracey, W. J. Trjitzinsky, A. W. Tucker, Bryant Tuckerman, J. W. Tukey, S. M. Ulam, G. B. Van Schaack, J. H. Van Vleck, H. E. Vaughan, Oswald Veblen, Abraham Wald, G. L. Walker, G. W. Walker, R. J. Walker, H. S. Wall, Charles Walmsley, J. L. Walsh, Warren Weaver, W. D. A. Westfall, A. P. Wheeler, H. S. White, A. L. Whiteman, P. M. Whitman, Hassler Whitney, J. K. Whittemore, G. T. Whyburn, W. M. Whyburn, D. V. Widder, Norbert Wiener, L. R. Wilcox, C. E. Wilder, S. S. Wilks, W. L. G. Williams, E. W. Wilson, Aurel Wintner, Audrey Wishard, E. E. Witmer, Jacob Wolfowitz, W. D. Wray.

The Colloquium Lectures on the subject *Analytic topology* were delivered by Professor G. T. Whyburn of the University of Virginia on Tuesday morning and afternoon and Wednesday and Thursday mornings. The presiding officers at the four lectures were, in order, President G. C. Evans, Professor J. R. Kline, Vice President C. R. Adams, and Vice President T. C. Fry.

On Thursday afternoon, by invitation of the Program Committee, Professor Leonard Carlitz of Duke University gave an address entitled *Arithmetic of polynomials in a Galois field.*

President G. C. Evans presided at the business meeting and Professor J. L. Walsh at the general session on Thursday morning; Professor Gordon Pall at the general session on Thursday afternoon. The presiding officers of the sectional sessions were as follows: Analysis, Professors L. W. Cohen, J. F. Randolph, I. M. Sheffer, Gabor Szegö; Algebra, Topology and Foundations, Dr. Marshall Hall; Algebra and Theory of Numbers, Professor B. W. Jones; Geometry, Professor M. G. Gaba; Mathematical Statistics (a joint session with the Institute of Mathematical Statistics), Professor B. H. Camp.

Headquarters for the meeting were in Thayer Hall. The dormitories of the college were available to the members of the three organizations and their guests.

Those attending the meetings were entertained Monday evening by a music recital, by moving pictures showing winter sports at Dartmouth, and by an exhibition of string models. On Monday after-

noon, tea was served at the Graduate Club by the ladies of the Dartmouth faculty.

A dinner for the members of the Society, the Association and the Institute was held Tuesday evening in Thayer Hall. Professor B. H. Brown was toastmaster. The speakers were Dean E. G. Bill, Dean R. G. D. Richardson, Professors P. A. Smith, Hassler Whitney, and S. S. Wilks. The attendance was 324.

Wednesday afternoon was devoted to an excursion to Franconia Notch.

The Bell Telephone Laboratories exhibited a machine for computing with complex numbers. The recording instrument at Hanover was connected by telegraph with the computing mechanism in New York. This machine was available to members from 11 A.M. to 2 P.M. each day of the meetings.

A resolution by Professor E. J. Purcell thanking the President, officers and members of the Mathematics Department of Dartmouth College for their competent arrangements and cordial hospitality, was adopted unanimously at the general session Thursday morning.

The Board of Trustees met on Monday at 6:00 P.M. in Thayer Hall. It reviewed the financial affairs of the Society with a view to a report to the Council.

There were three meetings of the Council, one being a joint session with the Board of Governors of the Mathematical Association of America. This joint session to hear the report of the War Preparedness Committee was held in Dartmouth Hall at 9:00 P.M. on Monday. The other meetings of the Council were in Thayer Hall at 4:30 P.M. on Tuesday and at 9:00 P.M. on Wednesday.

The Secretary announced the election of the following sixteen persons to membership in the Society:

Miss Helen C. Brodie, New York, N. Y.;
Professor D. van Dantzig, Technische Hoogeschool, Delft, Netherlands;
Professor Joseph J. Dennis, Clark University, Atlanta, Ga.;
Mr. Albert B. Farnell, University of Oklahoma;
Professor James Norman Goodier, Cornell University;
Dr. L. Louise Johnson, Reed College, Portland, Ore.;
Professor Eugene Leimanis, University of Latvia, Riga, Latvia;
Dr. Beppo Levi, The Mathematical Institute, Rosario, Argentina;
Dr. Harry Miller Lydenberg, New York, N. Y.;
Mr. (Albert) Wayne McGaughey, University of Cincinnati;
Sister Teresa M. Madden, College of Our Lady of the Elms, Chicopee, Mass.;
Mr. James R. Newman, New York, N. Y.;
Dr. Maxwell Reade, Ohio State University;
Dr. Charles V. Robinson, University of Missouri;

Professor Arthur Rosenthal, University of Michigan;
Dr. Walter Stephen Snyder, Ohio State University.

The following appointments by President Evans were reported: as representative of the Society at the dedication of buildings at the University of Colorado on June 8–9, Professor A. J. Kempner; as chairman of the Committee on Publicity (to replace Professor C. N. Moore, who resigned), Mr. R. M. Foster; as an additional member of this Committee, Professor J. B. Rosser (the newly constituted committee is as follows: Messrs. R. M. Foster (chairman), Saunders MacLane, G. B. Price, J. B. Rosser, and J. M. Thompson); as representative of the Society at the Conference on Science, Philosophy, and Religion and Their Relation to the Democratic Way of Life, held at the Jewish Theological Seminary of America in New York City on September 9–11, 1940, Professor Hermann Weyl; as representative of the Society at the inauguration of George N. Shuster as fifth president of Hunter College on October 8–11, Professor H. S. White.

Meetings of the Society during 1941 were set as follows: in New York City on February 22; in Stanford University, California on April 5; in Washington, D. C. on May 2–3; at the University of Chicago on September 2–6; in New York City on October 25. Summer Meetings were arranged for Cornell University in 1942 and for the University of Colorado in 1943.

An invitation to meet at the University of New Hampshire on June 23–27, 1941, in connection with the Congress of Sciences and Humanities and the American Association for the Advancement of Science was received.

The Committee on War Preparedness presented an extended report, which is summarized in an article by the Chairman, Professor Marston Morse, in the September number of this Bulletin. A Roster of Scientific Personnel is being compiled by the National Resources Planning Board of the U. S. Government, and Dean L. P. Eisenhart, Professor G. C. Evans, and Dean R. G. D. Richardson represent mathematics on the committee.

The Committee on Scientific Aids to Learning (supported by funds from the Carnegie Corporation) has appropriated a sum not to exceed $10,000 for microfilm reading machines, the distribution of which is to be for the benefit of Mathematical Reviews and to be arranged by the Executive Committee of that journal.

Reports were sent in July to the Carnegie Corporation and Rockefeller Foundation regarding their grants to Mathematical Reviews.

Certain invitations to give hour addresses were announced: Pro-

fessors J. B. Rosser and H. S. Wall for the November 1940 meeting in Detroit, Professors Saunders MacLane and Leo Zippin for the 1940 Annual Meeting, and Professor M. R. Hestenes for the 1941 April meeting in Chicago.

The sixteenth Josiah Willard Gibbs Lecture is to be given by Professor Sewall Wright at the Summer Meeting of 1941 in Chicago. The committee to select Gibbs lecturers for 1942 and 1943 consists of Professors H. P. Robertson (chairman), N. H. McCoy, and J. L. Synge.

Professors H. E. Buchanan and H. J. Ettlinger were appointed representatives on the Council of the American Association for the Advancement of Science for 1941 and Professor F. D. Murnaghan was appointed representative on the National Research Council for a period of three years beginning July, 1941.

The report of the Committee to Consider Possible Freer Methods of Nominating and Electing Council Members as adopted by the Council provides for a decrease in the number of ex-officio members of the Council, the opportunity for choice on the ballot in the election of vice presidents and elected members of the Council, and the chance on the ballot to suggest names for the various offices for the year following. A committee consisting of Professors W. M. Whyburn (chairman), Alonzo Church, and Philip Franklin was appointed to make appropriate revisions of the By-Laws with a view to presenting them for adoption by the Society. Provision is to be made on the Council for a representative of Mathematical Reviews.

A Committee on the Award of the Cole Prize in the Theory of Numbers for the period 1936–1940 was appointed, consisting of Professors H. H. Mitchell (chairman), A. J. Kempner, and Hermann Weyl.

A committee consisting of Professor Richard Courant (chairman), Mr. R. M. Foster, and Professor Harold Hotelling was named to consider the problem of holding informal discussion groups in regard to important papers read before the Society which touch on applications.

Titles and cross references to the abstracts of the papers read are given below. The papers were read as follows: papers 1–8 in the section for Analysis on Tuesday morning; papers 9–13 in the section for Geometry on Tuesday morning; papers 14–19 in the section for Mathematical Statistics on Tuesday afternoon; papers 20–23 in the section for Analysis on Tuesday afternoon; papers 24–31 in the section for Analysis on Wednesday morning; papers 32–38 in the section for Algebra, Topology and Foundations on Wednesday morning; papers 39–43 in the General Session on Thursday morning; papers 44–

50 in the section for Analysis on Thursday afternoon; papers 51–55 in the section for Algebra and Theory of Numbers on Thursday afternoon; and papers 56–96, whose abstract numbers are followed by the letter *t*, were read by title. Dr. Wundheiler was introduced by Professor D. J. Struik, Dr. Brown by Professor S. S. Wilks, Professor Zygmund by Professor T. R. Hollcroft, Dr. Hutchinson by Professor D. J. Struik, Professor Alexandroff by Professor P. A. Smith, Dr. Bergman by Dr. D. C. Spencer, Professor Hua by Professor T. R. Hollcroft, Dr. Manel by Professor Richard Courant, Professor Pólya by Professor Einar Hille, and Mr. Rosenthall by Professor E. T. Bell. Paper 1 was presented by Dr. Stephan Bergman, paper 4 by Mr. Epstein, paper 8 by Dr. Poritsky, paper 12 by Professor Struik, paper 21 by Dr. Boas, paper 22 by Dr. Halmos, paper 30 by Dr. Spencer, paper 31 by Professor Wiener, paper 38 by Dr. Hall, paper 39 by Professor Bochner, and paper 46 by Dr. Alaoglu.

1. Abe Gelbart: *On approximations of functions of two complex variables regular in multiply-connected four-dimensional domains.* (Abstract 46-9-418.)

2. K. O. Friedrichs: *On hyperbolic differential operators.* (Abstract 46-9-416.)

3. Orrin Frink: *Series expansions in linear vector space.* (Abstract 46-9-417.)

4. D. G. Bourgin and Benjamin Epstein: *A class of kernels generated by a Laplace-Mellin transformation.* (Abstract 46-11-448.)

5. Isaac Opatowski: *On the motion of an electric particle.* (Abstract 46-11-468.)

6. D. C. Harkin: *Fourier series as limits of Gaussian sums.* (Abstract 46-11-457.)

7. Einar Hille: *A class of differential operators of infinite order.* (Abstract 46-11-491.)

8. Hillel Poritsky and H. D. Snively: *Doubly periodic networks.* (Abstract 46-11-471.)

9. Alexander Wundheiler: *Some basic notions in meta-geometry.* (Abstract 46-9-445.)

10. Marshall Hall: *Projective planes.* (Abstract 46-9-421.)

11. W. L. G. Williams: *The application of hyperbolic complex numbers to the geometry of the triangle.* (Abstract 46-11-482.)

12. D. J. Struik and Alexander Wundheiler: *Affine representation of systems of forces.* (Abstract 46-9-439.)

13. E. J. Purcell: *Space Cremona transformations of order* $m+n-1$. (Abstract 46-9-433.)

14. Hilda P. Geiringer: *A generalization of the law of large numbers.* (Abstract 46-11-455.)

15. S. S. Wilks: *On the problem of two samples from normal populations with unequal variances.* (Abstract 46-11-481.)

16. Harold Hotelling: *Experimental determination of the maximum of an empirical function.* (Abstract 46-9-424.)

17. Abraham Wald: *Asymptotically shortest confidence intervals.* (Abstract 46-11-479.)

18. G. W. Brown: *Reduction of certain composite statistical hypotheses.* (Abstract 46-11-449.)

19. Jerzy Neyman: *Conception of equivalence in the limit of statistical tests and its application to certain forms of χ^2 and to the tests based on the λ principle.* (Abstract 46-11-467.)

20. J. F. Randolph: *Some properties of sets of the Cantor type.* (Abstract 46-11-472.)

21. R. P. Boas and D. V. Widder: *Functions with positive differences.* (Abstract 46-9-404.)

22. P. R. Halmos and R. A. Leibler: *Square roots of measure preserving transformations.* (Abstract 46-9-422.)

23. Antoni Zygmund: *Power series of the class H^λ on the circle of convergence.* (Abstract 46-11-483.)

24. J. G. Herriot: *Nörlund summability of double Fourier series.* Preliminary report. (Abstract 46-9-423.)

25. M. H. Heins: *A proof of Picard's theorem by the fundamental theorem on the conformal mapping of simply-connected Riemann surfaces.* Abstract 46–11-459.)

26. I. M. Sheffer: *On the singularities of functions under certain linear transformations.* (Abstract 46-11-475.)

27. I. J. Schoenberg: *On positive definite functions on spheres.* (Abstract 46-11-474.)

28. W. J. Trjitzinsky: *Properties of growth for solutions of differential equations of dynamical type.* (Abstract 46-11-478.)

29. J. M. H. Olmsted: *Lebesgue theory on a Boolean algebra.* Preliminary report. (Abstract 46-9-430.)

30. Stefan Bergman and D. C. Spencer: *On distortion in pseudo-conformal mapping.* (Abstract 46-9-402.)

31. Norbert Wiener and Aurel Wintner: *Discrete chaos.* (Abstract 46-11-480.)

32. P. M. Whitman: *Splittings of a lattice.* (Abstract 46-9-442.)

33. L. R. Wilcox: *A topology for semi-modular lattices.* (Abstract 46-9-443.)

34. O. G. Harrold (National Research Fellow): *Characterizations of some continua by continuous functions.* Preliminary report. (Abstract 46-11-458.)

35. Everett Pitcher: *The cap heights of a sum function on a product space.* (Abstract 46-11-470.)

36. S. C. Kleene: *Recursive predicates and quantifiers.* Preliminary report. (Abstract 46-11-464.)

37. Samuel Eilenberg: *On homotopy groups.* (Abstract 46-11-453.)

38. D. W. Hall and W. T. Puckett: *Strongly arcwise connected sets.* (Abstract 46-11-456.)

39. Salomon Bochner and I. J. Schoenberg: *On positive definite functions on compact spaces.* (Abstract 46-11-447.)

40. Gordon Pall: *On the arithmetic of ternary quadratic forms.* (Abstract 46-11-469.)

41. R. E. Langer: *On the theory of irregular differential boundary problems.* Preliminary report. (Abstract 46-11-465.)

42. G. C. Evans: *Surfaces of minimum capacity.* (Abstract 46-9-413.)

43. H. B. Curry: *The inconsistency of certain formal logics.* (Abstract 46-11-451.)

44. Warren Ambrose: *Change of velocities in a continuous ergodic flow.* Preliminary report. (Abstract 46-11-446.)

45. Benjamin Epstein: *A contribution to the theory of two complex variables in certain domains.* (Abstract 46-9-412.)

46. Leon Alaoglu, Garrett Birkhoff, and B. J. Pettis: *General ergodic theorems.* II. Preliminary report. (Abstract 46-9-394.)

47. Fritz John: *Discontinuous convex solutions of difference equations.* (Abstract 46-9-425.)

48. J. H. Curtiss: *Necessary conditions in the theory of interpolation in the complex domain.* (Abstract 46-7-378.)

49. Otto Szász: *On the jump of almost periodic functions and Fourier integrals.* (Abstract 46-9-399.)

50. H. T. Fleddermann: *The equality of measure functions.* (Abstract 46-9-414.)

51. L. C. Hutchinson: *On the classification of the trivector.* (Abstract 46-11-460.)

52. Mark Kac: *On a question of Ramanujan.* (Abstract 46-9-426.)

53. J. S. Frame: *The double cosets of a finite group.* (Abstract 46-9-415.)

54. A. T. Brauer: *On a property of k consecutive integers.* (Abstract 46-9-405.)

55. John Dyer-Bennet: *A note on partitions of the set of positive integers.* (Abstract 46-9-411.)

56. Paul Alexandroff: *General combinatorial topology.* (Abstract 46-7-374-*t*.)

57. Stefan Bergman: *Boundary values of functions satisfying partial differential equations of elliptic type.* (Abstract 46-9-401-*t*.)

58. R. P. Boas: *A general moment problem.* (Abstract 46-9-403-*t*.)

59. Herbert Busemann and Walther Mayer: *On the foundations of calculus of variations.* (Absract 46-9-406-*t*.)

60. Leonard Carlitz: *On the Staudt-Clausen theorem.* (Abstract 46-11-450-*t*.)

61. R. H. Cole: *The expansion problem associated with an ordinary linear differential equation and boundary conditions applying at a set of collinear points.* (Abstract 46-9-407-*t*.)

62. H. V. Craig: *On extensors.* (Abstract 46-9-408-*t*.)

63. H. B. Curry: *A revision of the fundamental rules of combinatory logic.* (Abstract 46-9-409-*t*.)

64. H. B. Curry: *Consistency and completeness of the theory of combinators.* (Abstract 46-9-410-*t*.)

65. M. M. Day: *Ordered sets and closures.* (Abstract 46-9-395-*t*.)

66. Samuel Eilenberg: *Local connectedness in higher dimensions.* (Abstract 46-11-452-*t*.)

67. C. H. Forsyth: *Statistical interpretations of the first auxiliaries of the Gaussian symmetrical method.* (Abstract 46-11-454-*t*.)

68. H. L. Garabedian, H. S. Wall, and Einar Hille: *Hausdorff means and integral equations.* (Abstract 46-9-397-*t*.)

69. L. J. Green: *Twisted cubics associated with a space curve.* II. (Abstract 46-9-419-*t*.)

70. D. W. Hall and W. T. Puckett: *Arc-preserving transformations of a certain class of spaces.* (Abstract 46-9-420-*t*.)

71. O. G. Harrold (National Research Fellow): *A note on strongly irreducible maps of an interval.* (Abstract 46-7-381-*t*.)

72. Loo-Keng Hua: *Some "Anzahl" theorems in the theory of groups of prime-power order.* (Abstract 46-11-492-*t*.)

73. Loo-Keng Hua: *The lattice points in a circle.* (Abstract 46-11-493-*t*.)

74. W. H. Ingram: *A generalization of Erhard Schmidt's solution of the nonhomogeneous integral equation.* Preliminary report. (Abstract 46-11-494-*t*.)

75. Nathan Jacobson: *Restricted Lie algebras of characteristic p.* (Abstract 46-11-461-*t*.)

76. R. L. Jeffery: *Integration in abstract space.* (Abstract 46-11-462-*t*.)

77. F. B. Jones: *Quasi-continuous curves and certain boundary problems.* (Abstract 46-11-463-*t*.)

78. Edward Kasner and J. J. DeCicco: *Equilong and conformal transformations of period two.* (Abstract 46-9-398-*t*.)

79. W. H. McEwen: *On the degree of convergence of a series associated with an integro-differential system.* (Abstract 46-9-427-*t*.)

80. W. G. Madow: *Contributions to the theory of the representative method of sampling.* (Abstract 46-11-466-*t*.)

81. Bella Manel: *Conformal mapping of multiply connected domains on the basis of Plateau's problem.* (Abstract 46-7-384-*t*.)

82. H. E. Newell: *On the asymptotic forms of the solutions of an ordinary linear matric differential equation in the complex domain.* Preliminary report. (Abstract 46-9-428-*t*.)

83. Rufus Oldenburger: *La teoria de los polinomios de orden superior.* (Abstract 46-9-429-*t*.)

84. F. W. Perkins: *Polygonal and polyhedral means of functions.* Preliminary report. (Abstract 46-9-431-*t*.)

85. G. Pólya: *Sur l'existence de fonctions entières satisfaisant à certaines conditions linéaires.* (Abstract 46-9-432-*t*.)

86. W. T. Reid: *A new class of self-adjoint boundary value problems.* (Abstact 46-11-473-*t*.)

87. J. F. Ritt: *On a type of algebraic differential manifold.* (Abstract 46-9-434-*t*.)

88. L. B. Robinson: *Singular solutions of nonlinear functional equations.* (Abstract 46-9-435-*t*.)

89. R. M. Robinson: *Stencils for solving* $x^2 \equiv a \pmod{m}$. (Abstract 46-9-436-*t*.)

90. E. Rosenthall: *On the diophantine equation* $x^3+y^3=z^3+w^3$. (Abstract 46-9-437-*t*.)

91. A. R. Schweitzer: *Concerning general abstract relational spaces.* II. (Abstract 46-9-438-*t*.)

92. D. C. Spencer: *On an inequality of Grunsky.* (Abstract 46-11-476-*t*.)

93. Gabor Szegö: *Power series with multiply monotonic sequences of coefficients.* (Abstract 46-11-477-*t*.)

94. J. L. Walsh and W. E. Sewell: *Degree of polynomial approximation to analytic functions. Problem* β. (Abstract 45-7-369-*t*.)

95. J. H. C. Whitehead: *On incidence matrices, nuclei and homotopy types.* (Abstract 46-9-441-*t*.)

96. Orla V. Wood: *On relations between solutions of the differential equation of the second order with four regular singular points.* (Abstract 46-9-444-*t*.)

T. R. Hollcroft,
Associate Secretary

BOOK REVIEWS

La Théorie des Spineurs. By Élie Cartan. (Actualités Scientifiques et Industrielles, nos. 643 and 701.) Paris, Hermann, 1938. 95 pp. and 91 pp.

The theory of spinors has been treated in the literature from three points of view: the infinitesimal, the algebraic and the geometric. Cartan originated the theory in its infinitesimal aspect in 1913 and he has now written a full length account in which it has been his intention "de développer systematiquement la théorie des spineurs en donnant de ces êtres mathématiques une définition purement géométrique."

Cartan's definition of a spinor (§§93 and 109) is based upon the one-to-two correspondence between the ν-vectors which are coordinates of linear spaces of ν dimensions on the fundamental quadric cone in 2ν (or $2\nu+1$) dimensions and certain vectors with 2^ν components, called "simple" spinors. Despite this geometric introduction to the concept of a spinor, knowledge of the algebraic theory of the representation of linear groups will be of more use to the reader than any acquaintance with the classical ideas of geometry.

The first two chapters of the first volume do not contain the word "spinor." Instead, they include an elementary discussion of the following topics: euclidean space (including the complex and the real definite and indefinite cases), rotations and reflections, multivectors, definition of tensors, tensor algebra, irreducible and reducible tensors, matrices (their algebra and a brief account of unitary, orthogonal and hermitian matrices), and the irreducibility of multivectors. Any reader already acquainted with the material in these forty-seven pages will delight in the concise elegance of the exposition while others will benefit from its lucidity and accuracy.

In the third and fourth chapters the spinor theory is developed in detail for a three dimensional space. The components of a spinor are first introduced as specific functions of the components of an isotropic vector, affected however by an ambiguous sign. After showing that a spinor is a "euclidean tensor," the author introduces in a natural fashion the relation $(x_1)^2+(x_2)^2+(x_3)^2=XX$, where X is a matrix with linear forms in x_1, x_2 and x_3 as elements and the left member is a scalar matrix. This association of vectors and matrices is used to obtain the double-valued representation of the rotation group by linear transformations on spinors. Chapter III concludes with a discussion of reality conditions in the euclidean and pseudo-euclidean

cases. In Chapter IV several theorems on the linear representations of the various rotation groups are proved by making use of the infinitesimal generators of the groups.

The first chapter of the second volume extends the spinor theory to spaces of any odd number of dimensions while the following chapter does the same for spaces of an even number of dimensions. In each of these cases the chapter closes with a brief discussion of reality conditions.

The last three chapters (totalling thirty pages) are entitled: Spinors in the Space of Special Relativity; Linear Representations of the Lorentz Group; and Spinors and Dirac Equations in Riemannian Geometry.

Comparison of this exposition of the spinor theory with the work of Veblen and his students[1] reveals a surprisingly small amount of common material. Thus Cartan does not employ Veblen's useful concept of a linear family of geometric transformations, which is a generalization of the well known linear family of involutions on the projective line. The avoidance of any use of the index notation is also interesting since the use of dotted indices has sometimes been regarded (quite erroneously) as a characteristic feature of the theory. These monographs do contain a full and excellent account of the group representation aspect of the theory and are an invaluable contribution to the literature of the subject.

Aside from a few misprints, the only error noted was on page 18 of the first volume, where the first lemma is false in case the fundamental quadratic form in n variables is of signature $\pm(n-2)$. Even in this case a weakened form of the lemma suffices for the following proofs.

WALLACE GIVENS

Gesammelte Werke. By Johannes Kepler. Vol. 1. *Mysterium Cosmographicum. De Stella Nova.* Edited by Max Caspar. 1938. 15+493 pp. Vol. 2. *Astronomiae Pars Optica.* Edited by F. Hammer. 1939. 467 pp. Vol. 3. *Astronomia Nova.* Edited by Max Caspar. 1937. 488 pp. Munich, Beck.

In 1936 Max Caspar published the *Bibliographia Kepleriana* as a part of this series.

The three imposing volumes of Kepler's collected works represent

[1] In his introduction Cartan refers to Veblen's work as unpublished, although in his bibliography he lists Veblen's paper in the Comptes Rendus Congrès Oslo and the reviewer's dissertation, which was done under Veblen's direction. The most complete publication of the work at Princeton was in the form of mimeographed notes on lectures by Veblen and Givens under the title "Geometry of Complex Domains."

in a way the first conclusive steps in an established mathematically harmonious universe. The desire for simplicity in the theory of the motions of the heavenly bodies activated both Copernicus and Kepler. The former attained some simplification but retained the difficult systems of epicycles by which Greek mathematicians had achieved an explanation of celestial phenomena sufficient for some eighteen hundred years of observation. The new path in theory taken by Copernicus was the result not so much of phenomena not explainable by the old theories but of the desire for a more evident mathematical simplicity.

Man's interest in the observed movements of the heavenly bodies is doubtless the most persistent intellectual occupation of record. The ancient Babylonians derived from these observations and records material for mathematical calculations. With the Greeks comes not only the use of the older observations and records but, further, the study of associated mathematical and physical problems, connected with light and theories of vision.

In India also the heavenly bodies secured intelligent attention, with simplified methods of computation, and with algebraic devices and the use of the half-chord and even the shadow function in astronomical problems.

All of these developments contributed to the Moslem interest in astronomy, geography, and trigonometry, and the material was communicated through their works and instruments to Europe in the twelfth to fifteenth centuries. Kepler, more than any other of the European group which includes also Copernicus, Galilei, Tycho Brahe and Newton, appreciated most explicitly the achievements of the past centuries and their manifold contributions towards new theories.

Kepler began his search for a harmonious universe with an inspiration which today would be termed a delusion. However, Kepler kept faith with the observations as recorded and the continuance of this faith led him to observations not reconcilable with his inspiration nor with the theories of Ptolemy or Copernicus. The conclusive observations were furnished by Tycho Brahe, with a few by Kepler himself, and the mathematical universe involved in Kepler's laws of planetary motions gradually evolved as a result of enormous computations, treated with profound mathematical insight, establishing definitely the orbits of the planets as ellipses.

Kepler began his interest in astronomy at the University of Tübingen. The struggle for an existence led him to the casting of horoscopes and to the publication of almanacs, which were largely

for such uses. Despite this occupation the scientist Kepler sought a mathematically harmonious universe. It seemed to him to come as an inspired thought "on July 19, 1595," as he records in a letter, that there were six planets and five regular solids, and that, in consequence, the latter could be circumscribed with the sun as center about the orbits of the planets and thus harmoniously, possibly, fill in the gaps between the six planetary orbits. In the *Mysterium Cosmographicum* of 1596 Kepler elaborates this thesis, arriving at only partially successful conclusions.

In 1594 when Kepler began to teach mathematics the Copernican theory was not yet widely accepted. In his study of astronomy under an ardent Copernican, M. Mästlin, at Tübingen, Kepler had acquired an insight into the problems, leading him to desire further information concerning the size and the orbits of the planets. As the varying speeds of the planets were known and their approximate relative distances from the sun, Kepler was able to attain preliminary success in circumscribing around the recorded orbits the five regular solids, "limited by the unchangeable laws of geometry." With these five solids the sphere as a sixth solid was employed for the outermost sphere of the fixed stars.

The haste with which Kepler composed his *Mysterium Cosmographicum* is explained by the author's impending marriage and the necessity for impressing the University officials of Graz where he was employed. Doubtless had Kepler lingered, as he did with later projects, the *Mysterium* might, indeed, have never seen the light. Kepler spent some seven months in Tübingen, on vacation from Graz, completing the manuscript and arranging for the supervision of the printing by his former teacher, Mästlin.

A friendly reception was accorded the *Mysterium Cosmographicum* despite the assumption of the Copernican theory and the peculiarities inherent in the Keplerian hypothesis.

In placing the sun at the center of the universe, Kepler arrived at an appreciation, often reiterated in his later works, of the sun as the center of force of our universe. Copernicus also refers to the sun as the center of power, but he actually used in his epicycles the centers of the orbits as centers; Kepler made his measurements as far as possible directly from the sun. In Chapter XX (p. 70) Kepler refers to the sun as not only the fountain head of light, but the source of life, movement and active force (*anima*); in a later edition Kepler uses the word "*vis*."

Naturally the completed *Mysterium Cosmographicum* of 1595 was sent to the most distinguished astronomer then living, Tycho Brahe,

and the later association of Kepler with Brahe hinged upon this connection. The scholarly documentation of an untenable theory established Kepler's fame as an astronomer of great ability. For more correct planetary orbits Tycho Brahe had the data, and it was with an eye to these that Kepler made contact with the great Danish observer of the heavens.

With the *Mysterium Cosmographicum* Kepler included the *Narratio Prima* which had been published in 1540 by G. J. Rheticus,[1] as a preliminary statement concerning the Copernican system. To this was added an appendix by M. Mästlin, *De dimensione orbium et sphaerarum coelestium.*

The modern editors have included in this volume with the *Mysterium* four of Kepler's minor astronomical treatises, published under the title, *De Stella Nova . . .*, 1606; separate title pages are included for the other three treatises. These were publications designed in large part to take advantage, financially, of somewhat wide popular interest in astronomical occurrences.

Late in 1599 Tycho Brahe invited Kepler to visit him in Prague. In February of 1600 Kepler began somewhat tentatively employment at the Observatory in Prague.

When Kepler first approached Tycho Brahe he rather naively supposed that Tycho would be able to hand to him complete data of the orbits of the planets, so that within two or three weeks Kepler himself could check the results with his theory. In fact it took Kepler years to compute the orbit of Mars, the first one to be completed.

Tycho Brahe was having his difficulties in 1599 when he moved to Prague with the mathematical computations in connection with his observations and particularly with the orbit of Mars. Shortly thereafter Kepler was practically without a job, and as Tycho Brahe needed a mathematical astronomer, Kepler found himself in 1600 in Prague as an associate with Tycho Brahe, replacing the astronomer Longomontanus, who returned to Denmark. The observations of Mars and the problems of their reduction were placed in the hands of Kepler.

While Kepler was occupied with these computations, Tycho Brahe died October 24, 1601. Kepler was for a time prevented from free use of the observations of Tycho Brahe. Partly because of these hindrances the complete construction of the orbit of Mars was not effected until some years after the death of Tycho and then even further delayed in publication.

[1] See Edward Rosen, *Three Copernican Treatises*, New York, Columbia University Press, 1939; the English translation of the Rheticus treatise is given.

In the interval Kepler attacked a number of problems in optics and in observational astronomy which were essential to any correct interpretation of the observations. Kepler's treatise on optics started with the work of Witelo as a basis. Kepler established a modern point of view for the science and even touched some of the problems shortly to be created by the introduction of the telescope.

The *Astronomiae Pars Optica* includes a wide range of discussions on the nature and properties of light, the functioning of the eye, the optical image, reflection and refraction of light rays, and astronomical optics including the lights and shadows of heavenly bodies as well as problems of parallax. Through the work of Witelo written about 1270 A.D. Kepler takes account of the work of the Arabs, notably Al-Hazan (Ibn al-Haitam) and the optics of Euclid and Ptolemy; the whole science was modernized and innumerable mistaken points of view eliminated.

Notably in connection with the observation of eclipses Kepler introduced quite new methods and hurried the book through the press in order to make his methods available to astronomers for an eclipse to occur on October 2 (12) of 1605.

Within six years after this work by Kepler appeared, observational astronomy was revolutionized by the telescope. Kepler was able to make more obvious contributions to the new theory, particularly of lenses, in his *Dioptrice* of 1611. The *Optics* of 1604 revived the problems of refraction and stimulated the definitive solution by Snell and Descartes.

The orbit of Mars had resisted the efforts of Tycho Brahe and Longomontanus. To this problem Kepler gave himself with utmost devotion over a period of five years. About Easter of 1605 Kepler abandoned the eccentric circles and the ovals to make an attempt at the ellipse as the orbit. In view of the attention given in the *Pars Optica* to the focal properties of the conics and their possible applications, one is justified in the conclusion that this study paved the way to Kepler's Laws of Planetary Motion.

While in 1601 Kepler had arrived at the orbit of Mars as in a plane inclined to the ecliptic, not until October 11 of 1605 did Kepler announce the path of Mars as an ellipse; this was done in a letter to the astronomer, Fabricius. This achievement more than any other marks the beginning of the modern theory of the movements of the heavenly bodies.

In a measure the crucial step taken by Kepler was to take the sun itself and not the center of the orbit of a planet about the sun as the center from which computations were made. In the Copernican the-

ory the center of the earth's orbit about the sun is taken as the center of the universe. Kepler's innovation moved the center (focus) to the sun itself and gave a mathematical law for the varying velocity of each planet in its orbit. Vital in the computations was new light upon the earth's orbit. In fact, the second law of Kepler that equal areas are swept out in equal times was first demonstrated by Kepler for the earth; the formulation gave the velocity as inversely proportional to the distance from the sun, the "radius-theorem." The final establishment for the orbit of Mars was dependent upon the corrections made possible by the revisions made in the earth's orbit.

In preparing for publication of Kepler's collected works translations proved to be out of the question; the editors have introduced the device of admirable summaries of contents of each volume. In addition there are notes, often giving an indication of the method of work of the author, and the genesis of the ideas; in fact, Kepler himself gives much information along this line, including even material finally rejected. The editors have also included in each volume numerous notes and for the periods concerned available lists of Kepler's correspondence, largely with brief statements of contents of the letters.

In every way the series, *Johannes Kepler Gesammelte Werke*, can be commended as worthy of the great genius of Kepler.

LOUIS C. KARPINSKI

An Introduction to the Theory of Functions of a Real Variable. By S. Verblunsky. Oxford, Clarendon Press, 1939. 11+169 pp. $4.25.

This text for students and teachers was written for the special purpose of furnishing a more rigorous and accurate treatment of the elements of the theory of functions of a real variable. It is based on notes of lectures delivered by Verblunsky to students in their first year at the University of Manchester. The subject matter is entirely standard, but the treatment involves much that is new and original—ingenious and elegant proofs for certain theorems and new approaches to some parts of the subject. The dominant feature of the book is the presentation of the subject as a body of deductions from specified hypotheses.

The material treated will be sufficiently indicated by the following list of chapter headings: Chapter I, Number; II, Sets and Functions; III, Convergence; IV, Continuity and the Derivative; V, The Elementary Functions; VI, Primitives; VII, Limits and Higher Derivatives; VIII, Integrals; IX, Series. The elementary character of the book should be noted. There is no treatment of the properties of sets

of points, of measure, of Lebesgue integration, or of other modern topics.

The treatment is different or unusual in a number of respects; some of them are the following. (a) In Chapter I there is a set of postulates from which all the results in the book are derived. These postulates do not form a mutually independent set; they were chosen in an effort to make the initial progress of the student rapid and easy. (b) In Chapter I also the principle of induction is stated as a postulate; it is then used constantly, and whenever occasion demands, in proofs throughout the book. (c) Both Heine's and Cauchy's definitions of continuity are given, but the treatment is based largely on the former. Although there is a proof that the two definitions are equivalent, attention is not called to the use in it of the axiom of choice. Two definitions of a continuous function $f(x)$ are given, one for a closed interval (Chapter IV, p. 51) and another for an open interval (Chapter VI, p. 83). Finally, the treatment of continuity is separated from the treatment of limits of functions of a continuous variable, the former being placed first in Chapter IV and the latter afterward in Chapter VII. (d) The treatment of the derivative, placed as it is in Chapter IV before the study of limits in Chapter VII, obviously involves new features. The derivative of $f(x)$ at $x=\xi$ is shown to be the value $\phi(\xi)$ of an auxiliary function $\phi(x)$ which is continuous at $x=\xi$. There follow elegant derivations of all the standard properties of the derivative. (e) The treatment of the elementary functions (exponential, logarithmic, trigonometric, hyperbolic) in the short space of fourteen pages in Chapter V is a gem. (f) The introduction of an equivalence relation indicates the care used in the treatment of primitives in Chapter VI. The familiar theorem now reads: a primitive of the sum of two functions is equivalent (not *equal*!) to the sum of primitives of these functions. (g) The integral treated in Chapter VIII is essentially a special case of the Perron integral. It is quite general and integrates with equal ease bounded functions and certain functions which become infinite at a finite number of points. It is shown that for a continuous function it is possible to approximate to the integral by the Riemann sums; in general, however, there is no treatment of integrals as the limits of sums. (h) The book contains a large number of examples. Some illustrate definitions and theorems, and others are additional propositions which are essential for later developments. Most of the examples are accompanied by proofs.

To this reviewer it seems that Verblunsky has succeeded admirably in his effort to write a rigorous and accurate introduction to the theory of functions of a real variable. Furthermore, the literature has been

enriched permanently by original treatments of certain topics and the elegant proofs of numerous theorems which he has contributed. At the same time, the book has certain characteristics which this reviewer finds unfortunate in a text for beginners. They are these. (a) The subject has been completely divorced from its intuitive background and also from its historical development. The book contains no figures, and there is no indication of the origin of any of the ideas involved or of their applications. There are no references to the history of the subject. Even the names of mathematicians have been omitted; those of Rolle and MacLaurin are the only ones this reviewer could find. (b) There is an authoritative air of finality about the book. There is no suggestion that other treatments can be given for many of the topics, or that interesting extensions and generalizations are possible in some cases. A book with these characteristics is not likely to offer much *inspiration* to a beginning student. Nevertheless, it is a valuable text for superior students who are reading independently and for classes of an inspiring lecturer.

The printing has been done with an accuracy worthy of the author's success in producing a book of high mathematical quality. The reviewer noticed only three typographical errors, all of which were small and obvious; no one would cause confusion even to the beginning student. The author and printer are to be congratulated on producing an excellent mathematical text.

G. Baley Price

NOTES

The University of Pennsylvania celebrated the two hundredth anniversary of its origin in September 1940. Professor Hermann Weyl of the Institute for Advanced Study gave an address on *The mathematical way of thinking*. Dr. T. C. Fry of the Bell Telephone Laboratories was chairman of the symposium on statistical methods in engineering. Associate Professor S. S. Wilks of Princeton University and Dr. W. A. Shewhart of the Bell Telephone Laboratories participated in this symposium.

The following members of the American Mathematical Society are in residence at the Institute for Advanced Study for all or part of this academic year: Dr. Warren Ambrose, Dr. Valentin Bargmann, Assistant Professor D. G. Bourgin of the University of Illinois, Dr. A. T. Brauer, Professor Orrin Frink of Pennsylvania State College, Dr. Guido Fubini, Dr. Kurt Gödel, Dr. P. R. Halmos, Dr. M. H. Heins, Dr. Dorothy Maharam, Assistant Professor Gordon Pall of McGill University, Dr. Abraham Schwartz, Dr. C. E. Shannon, Dr. Seymour Sherman, Assistant Professor A. H. Taub of the University of Washington, Dr. R. M. Thrall, and Professor W. J. Trjitzinsky of the University of Illinois.

The following persons are also to be at the Institute for Advanced Study during a part or all of the current academic year: Dr. C. L. Critchfield, Professor Shizuo Kakutani of Osaka Imperial University, Professor Wolfgang Pauli of Zürich Polytechnicum, and Professor C. L. Siegel of the University of Göttingen.

Assistant Professor R. G. Sanger of the University of Chicago has been awarded one of three $1000 awards given at the University of Chicago to selected teachers for excellence in undergraduate teaching.

Professor H. H. Dalaker of the University of Minnesota has been given the title emeritus.

Professor Hermann Weyl of the Institute for Advanced Study has been awarded an honorary doctorate of science by the University of Pennsylvania.

Dr. Frances E. Baker of Mount Holyoke College has been promoted to an assistant professorship.

Dr. E. F. Beckenbach of Rice Institute has been appointed to an assistant professorship at the University of Michigan.

Dr. J. H. Curtiss of Cornell University has been promoted to an assistant professorship.

Dr. Churchill Eisenhart of the University of Wisconsin has been promoted to an assistant professorship.

Professor W. B. Ford of the University of Michigan has retired.

Professor E. G. Harrell of the North Dakota State Teachers College, has been appointed to a professorship at the State Teachers College, Platteville, Wisconsin.

Mr. L. B. Hedge of Brown University has been appointed to an assistant professorship at The Citadel.

Dr. C. C. Hurd of Michigan State College has been promoted to an assistant professorship.

Dr. J. F. Kubis of Fordham University has been promoted to an assistant professorship.

Dr. S. B. Littauer of the U. S. Naval Academy at Annapolis has been promoted to an assistant professorship.

Professor C. C. MacDuffee of the University of Wisconsin has been appointed to a professorship at Hunter College.

Dr. W. H. Myers of the University of Utah has been appointed to an assistant professorship at San Jose Junior College, San Jose, California.

Assistant Professor N. N. Royall of The Citadel has been appointed to an associate professorship at Winthrop College, Rock Hill, South Carolina.

Dr. S. S. Smith of the University of Utah has been promoted to an assistant professorship.

Associate Professor R. C. Stephens of Knox College has been promoted to a professorship.

Dr. J. A. Ward of the Tennessee Polytechnic Institute has been appointed to an associate professorhip at Delta State Teachers College, Cleveland, Mississippi.

Dr. L. R. Wilcox of the University of Wisconsin has been appointed to an assistant professorship at the Illinois Institute of Technology.

Associate Professor P. A. De Vore of Central Missouri State Teachers College has been granted a leave of absence of this academic year.

Dr. R. H. Fox of the Institute for Advanced Study has been made an associate at the University of Illinois.

The following appointments to instructorships are announced: Brooklyn College: Mr. Saul Gorn, Dr. D. C. Harkin; College of the City of New York: Dr. G. N. Garrison; Cooper Union: Dr. Alvin Sugar; Duke University: Miss Mary E. Layne; Eastern Washington College of Education, Cheney, Washington: Mr. R. F. Bell; Gogebic Junior College, Ironwood, Michigan: Mr. K. F. McLaughlin; Illinois Institute of Technology: Dr. Herbert Busemann, Dr. J. J. DeCicco; Iowa State College: Dr. A. T. Lonseth; Johns Hopkins University: Dr. R. B. Kershner; University of Michigan: Dr. G. E. Hay, Dr. Wilfred Kaplan; University of Minnesota: Dr. J. M. H. Olmsted; University of Missouri: Dr. M. E. Shanks, Dr. J. V. Wehausen; University of Nevada: Mr. E. M. Beesley; North Carolina State College: Dr. Abe Gelbart; Oklahoma Agricultural and Mechanical College: Dr. P. E. Lewis; Purdue University: Mr. C. D. Olds; Southern Methodist University: Dr. D. W. Starr; Texas Technological College: Dr. P. W. Gilbert; Vanderbilt University: Dr. John Dyer-Bennet; University of Wisconsin: Dr. S. M. Ulam.

Mr. Jonathan Hodgkinson of Jesus College, Oxford University, died June 21, 1940, at the age of fifty-four years.

The death of Professor A. E. H. Love of Oxford University has been reported.

Dr. Frederic Marty lost his life June 14, 1940 while in the service of France.

Dr. E. G. C. Poole of Oxford University died June 28, 1940, at the age of forty-nine years.

Dean F. G. Wren of Tufts College died July 17, 1940, at the age of sixty-six years. He had been a member of the Society since 1922.

ABSTRACTS OF PAPERS

SUBMITTED FOR PRESENTATION TO THE SOCIETY

The following papers have been submitted to the Secretary and the Associate Secretaries of the Society for presentation at meetings of the Society. They are numbered serially throughout this volume. Cross references to them in the reports of the meetings will give the number of this volume, the number of this issue, and the serial number of the abstract.

446. Warren Ambrose: *Change of velocities in a continuous ergodic flow.* Preliminary report.

A flow is a 1-parameter group T_t, $-\infty < t < \infty$, of measure preserving transformations of a space into itself. It is measurable (continuous) if T_tP is a measurable (continuous) (P, t)-function. If S is any measure preserving transformation of a space Ω into itself, then a flow can be "built on S" as follows: consider the product space of Ω with $0 \leqq x < 1$, with measure defined multiplicatively. Then define the flow by $T_t(P, x) = (S^nP, t+x-n)$, where n is equal to the integral part of $t+x$. It is shown that for any continuous ergodic flow on a separable metric space of finite measure the velocities along the streamlines can be altered to obtain a flow built on a measure preserving transformation, that is, a subspace of the original space can be found whose product with $0 \leqq x < 1$ is, with respect to measure properties, the original space (where measure on that original space is now the measure invariant under the altered flow) and such that the altered flow is built on a measure preserving transformation on this subspace. It is intended to use this theorem in a study of spectral properties which are invariant under a change of velocities. (Received August 5, 1940.)

447. Salomon Bochner and I. J. Schoenberg: *On positive definite functions on compact spaces.*

The expansion theorem for positive definite functions on finite-dimensional euclidean spheres, as stated in *On positive definite functions on spheres* by I. J. Schoenberg (abstract 46-11-474), is contained in a general theorem concerning such functions on compact spaces on which a transitive group of transformations is defined. In the general case, as in the special case, the function is given as a function of two independent points which in addition to being positive definite is invariant under simultaneous transformations of both points by the same group element. (Received August 5, 1940.)

448. D. G. Bourgin and Benjamin Epstein: *A class of kernels generated by a Laplace-Mellin transformation.*

The kernels discussed include those of the type of the generalized zeta function. The authors treat the inversion in L_2 of integrals with these kernels. (Received August 5, 1940.)

449. G. W. Brown: *Reduction of certain composite statistical hypotheses.*

The results obtained make it possible to reduce a large class of "composite" statis-

tical hypotheses to equivalent "simple" hypotheses. The fundamental theorem established states essentially that if two distributions give rise, in sampling, to the same distribution of the set of differences between observations, then one distribution must be a translation of the other, subject to a condition requiring that the characteristic function of one of the distributions be such that any interior intervals of zeros be not too large. The result is established by means of the functional equation $\phi(t_1)\phi(t_2)\phi(-t_1-t_2)=\psi(t_1)\psi(t_2)\psi(-t_1-t_2)$ relating the characteristic functions. Similar results are obtained for scale, and combination of location and scale, and the corresponding situations in multivariate distributions. This type of uniqueness theorem permits one to reduce a composite statistical hypothesis involving an unknown location parameter (or scale, or both) to an equivalent simple hypothesis. (Received August 2, 1940.)

450. Leonard Carlitz: *On the Staudt-Clausen theorem.*

The principal object of this note is to simplify both the proof and formulation of the "polynomial" analogue of the Staudt-Clausen theorem given some time ago (Duke Mathematical Journal, vol. 3 (1937), pp. 503–517). (Received August 6, 1940.)

451. H. B. Curry: *The inconsistency of certain formal logics.*

In 1935 Kleene and Rosser published a proof that certain systems of formal logic were inconsistent in the sense that every formula which could be formulated in them was also demonstrable. This proof has been revised and simplified by the present author, who showed that the inconsistency depends, essentially, on the conjunction of two incompatible properties, called combinatorial and deductive completeness respectively (see abstract 43-3-118). The argument, in both the original proof and the revision, was a refinement of the Richard paradox. In the present paper it is shown that a contradiction can be derived by a much simpler argument based on the paradox of Epimenides. (Received August 6, 1940.)

452. Samuel Eilenberg: *Local connectedness in higher dimensions.*

A metric space Y which is locally p-connected in the homotopy sense for $k\leqq p\leqq n$ (notation: Y is LC_k^n) has the following property (P): Given any metric separable space X and a closed subset X_1 which is LC_0^{k-1} such that $\dim X-X_1\leqq n+1$, every continuous mapping $f(X_1)\subset Y$ has an extension $f(U)<Y$ where U is some open set containing X_1. If Y is locally 0-connected then the property (P) is equivalent with Y being LC_k^n. If the property (P) holds without the restriction $\dim X-X_1\leqq n+1$ then Y is said to be an absolute neighborhood retract starting from the dimension k (notation: Y is ANR_k). If Y is compact then Y is an absolute neighborhood retract if and only if Y is LC_0^{k-1} and ANR_k. A compact space Y which is ANR_1 may not be locally p-connected for any $p\geqq 0$; however, if Y is locally 0-connected then it is LC_0^n for every n. (Received August 9, 1940.)

453. Samuel Eilenberg: *On homotopy groups.*

Let Y be an arcwise connected topological space. A singular simplex in Y is a couple (G^p, f) where G^p is a closed euclidean p-simplex and f is a mapping $f(G^p)\subset Y$. After a proper equivalence relation has been introduced, the totality of all the singular cells in Y was shown by Lefschetz to form a closure-finite complex $S(Y)$. The nth homology group $H^n(S(Y))$ coincides with the homology group $H^n(Y)$ obtained using singular chains and cycles. Given $y_0\varepsilon Y$ and $m\geqq 0$ let $S_m(Y)$ be the subcomplex of $S(Y)$ consisting of all singular simplexes (G^p, f) such that every face of G^p of dimension less than

m is mapped by f into y_0. For every n the groups $H^{n,m}(Y)=H^n(S_m(Y))$ are considered where $0\leqq m\leqq n$. The groups $H^{n,0}(Y)$, $H^{n,1}(Y)$ and $H^n(Y)$ are isomorphic. If $n>1$ and the coefficients are integers, then $H^{n,n}(Y)$ is the nth homotopy group $\pi_n(Y)$ of Y. (Received August 9, 1940.)

454. C. H. Forsyth: *Statistical interpretations of the first auxiliaries of the Gaussian symmetrical method.*

While the Gaussian symmetrical method of applying least squares is well known and fully treated in mathematical literature, the fact that the first auxiliaries of that method form a veritable treasure house for most of the fundamental formulas of statistical theory (including those of means, dispersions and even of correlation coefficients) has not, apparently, been duly appreciated. The author applies the notation peculiar to statistical theory to discover all these formulas. (Received August 7, 1940.)

455. Hilda P. Geiringer: *A generalization of the law of large numbers.*

Let $V_1(x)$, $V_2(x)$, $\cdots$, $V_n(x)$ be n probability distributions which are not supposed to be independent and let $F(x_1, \cdots, x_n)$ be a "statistical function" of n observations in the sense of von Mises, $V_i(x)$, $i=1, \cdots, n$, indicating as usual the probability of getting a result equal to or less than x at the ith observation. Then it can be proved that under fairly general conditions $F(x_1, \cdots, x_n)$ converges stochastically to its theoretical value; or, in other words, that under these general conditions a great class of statistics $F(x_1, \cdots, x_n)$ are consistent in the sense of R. A. Fisher. Well known particular cases of this theorem result (a) if $F(x_1, \cdots, x_n)$ is taken for the average $(x_1+x_2+\cdots+x_n)/n$ of the n observations, (b) if it is assumed that the $V_i(x)$ are independent distributions. (Received August 5, 1940.)

456. D. W. Hall and W. T. Puckett: *Strongly arcwise connected sets.*

In this paper two characterizations of strongly arcwise connected (see W. T. Puckett, this Bulletin, abstract 46-5-316) sets are obtained. The first characterization states that a locally connected continuum A is strongly arcwise connected if and only if every infinite collection of open sets in A contains an infinite subcollection intersecting some arc of A. Strong arcwise connectivity of a set at a point p is defined and a necessary and sufficient condition that the locally connected continuum M fail to be strongly arcwise connected at a point p is obtained. From this criterion it is deduced that an arbitrary locally connected continuum M is strongly arcwise connected at all save possibly a countable number of its points. If no two points separate M, then M is strongly arcwise connected. (Received September 11, 1940.)

457. D. C. Harkin: *Fourier series as limits of Gaussian sums.*

By means of the cyclotomic algebra which Gauss introduces in the seventh section of the *Disquisitiones Arithmeticae*, Fourier series can be had as limiting cases of sums similar to those there considered. (Received August 15, 1940.)

458. O. G. Harrold (National Research Fellow): *Characterizations of some continua by continuous functions.* Preliminary report.

In this paper a study is made of mappings of a given continuum into the interval (0, 1) which are characteristic. This method has been used by Mazurkiewicz, Whyburn

and Eilenberg. New characterizations are given for the continua of finite degree and the continua having no continuum of condensation. For the last mentioned class of continua, a subclass of all continuous mappings is found with respect to which this type of continuum has much the same properties as the class of all continuous mappings with respect to the class of all Peano continua. If a mapping of A into B is called of *almost all finite sections* provided that $f^{-1}(y)$ is finite, $y \in B$, except for a countable set of points each of which has a countable inverse (A.F.S.), then a necessary and sufficient condition that a Peano continuum admit an A.F.S. map into (0, 1) is that each dendrite in the continuum have this property. This is related to a problem proposed by Cech (Fundamenta Mathematicae, vol. 18). (Received August 6, 1940.)

459. M. H. Heins: *A proof of Picard's theorem by the fundamental theorem on the conformal mapping of simpy-connected Riemann surfaces.*

Let G_w^* denote the universal covering surface of G_w, the extended w-plane with three distinct points a, b, c deleted. G_w^* is simply-connected, and, as the universal covering surface of the Riemann surface G_w, G_w^* may be taken to be a Riemann surface. Hence G_w^* can be mapped one-to-one and conformally onto one and only one of the following canonical domains: (1) the extended z-plane, (2) the finite z-plane, (3) the interior of the unit circle in the z-plane. The first possibility is readily excluded. As for the second, let $w=w(z)$ denote a mapping function which allegedly maps the finite z-plane one-to-one and conformally onto G_w^*. $w=w(z)$ would be automorphic under a group of linear transformations mapping the finite z-plane onto itself. From this one infers readily that $w(z)$ must be doubly-periodic, simply-periodic, or a constant. In each case a contradiction is manifest. Hence G_w^* must be of hyperbolic type. Picard's theorem that a function, not a constant, meromorphic in the finite plane, can omit at most two values follows immediately. (Received August 6, 1940.)

460. L. C. Hutchinson: *On the classification of the trivector.*

In this paper the author finds by a new method the known canonical forms for trivectors, or alternating tensors of the third valency, in eight dimensions, as well as new forms which do not exist as canonical forms except in the real domain, and examines some of their geometrical properties. This paper is based on the author's work for the doctorate at the Massachusetts Institute of Technology. (Received August 6, 1940.)

461. Nathan Jacobson: *Restricted Lie algebras of characteristic p.*

In this paper certain identities previously noted connecting pth powers, $a+b$, $a\alpha$, α in the field and $[a, b]=ab-ba$ in an associative algebra of characteristic $p\neq 0$, are proved to be characteristic of these operations. This is shown by defining the concept of an abstract restricted Lie algebra and proving that any such algebra L may be obtained as a subset of an associative algebra closed under the above operations. Among the minimal associative algebras determined by L in this way, there is one which is universal in a certain sense. The derivation algebra of any algebra of characteristic p is an instance of a restricted Lie algebra. A notion of a restricted derivation of a restricted Lie algebra is defined. These form a restricted Lie algebra and may be used to define a restricted holomorph. If L has a finite basis it has a (1-1) representation by finite matrices. An analogue of Engel's theorem holds in this case. (Received August 9, 1940.)

462. R. L. Jeffrey: *Integration in abstract space.*

A theory of integration in a complete normed vector space is given which is shown to be equivalent to that given by G. Birkhoff (Transactions of this Society, vol. 38 (1935), pp. 357–378). This theory is in the spirit of the classical theories for real and complex variables. In particular it makes no use of convex sets, and, as a consequence of this, the developments are as simple as those of the classical theories. (Received August 3, 1940.)

463. F. B. Jones: *Quasi-continuous curves and certain boundary problems.*

A continuum M is said to be *quasi-connected at* P if P belongs to M and for each point X of $M-P$ there exists a continuum which lies in $M-X$ and contains an open subset of M containing P. A continuum which is quasi-connected at each of its points is said to be a *quasi-continuous curve.* The class of quasi-continuous curves includes as proper subclasses both the class of continuous curves and the class of semi-locally connected continua, whereas neither of the latter classes includes the other. The first part of the paper deals largely with the similarity between quasi-continuous curves and continuous curves in a Hausdorff space. The second part of the paper deals with boundary problems in a space whose only two-dimensional feature is the Jordan curve theorem. If P is a point of a continuum K in such a space so that (1) K is locally compact at P and (2) K is quasiconnected at P, then K together with all but a finite num ber of its complementary domains is connected im kleinen at P. As an interesting application of this theorem, it is shown that a proposition analogous to Whyburn's generalization of the Torhorst theorem for the plane holds true in these abstract spaces. (Received August 10, 1940.)

464. S. C. Kleene: *Recursive predicates and quantifiers.* Preliminary report.

Consider the following forms of expression: $R(n)$, $(Ex)R(n, x)$, $(x)R(n, x)$, $(x)(Ey)R(n, x, y)$, $(Ex)(y)R(n, x, y)$, $\cdots$. The variables $n, x, y, \cdots$ range over non-negative integers. The quantifiers $(E.)$ and $(.)$ express "there exists an . such that" and "for .," respectively. Choose one of the forms, and let R be a predicate (that is, propositional function) of the variables used in that form. Then the expression represents a predicate P of the variable n. As defined by Herbrand and Gödel, a general recursive predicate has a certain kind of algorithm, which, for given values of the variables, leads to a decision as to the truth or falsity in a finite number of steps. According to Church, this kind of algorithm is the most general possible. To each of the above forms, a general recursive R can be chosen, such that the resulting P is not given by any of the other forms with the same or fewer quantifiers, for any general recursive R whatsoever. This includes, for the first of the above forms, Church's result on an unsolvable problem of elementary number theory, and for the second, a part of Gödel's results on the incompleteness of formal deductive systems. The R chosen for each form is primitive recursive. (Received August 7, 1940.)

465. R. E. Langer: *On the theory of irregular differential boundary problems.* Preliminary report.

A matrix differential boundary problem $Y'=(\lambda R+B)Y$, $W_a Y(a)+W_b Y(b)=0$, is familiarly classifiable as regular, mildly irregular, or highly irregular, according

as the characteristic equation possesses or does not possess certain structural peculiarities. In any event, if an infinity of characteristic values exists, an "arbitrary" function may be formally associated with an "expansion" in characteristic solutions of the problem. Under appropriate conditions this expansion is known to be convergent or summable to the given function if the problem is respectively regular or mildly irregular. There is, however, little in the way of general theory for the highly irregular cases. This paper seeks to obtain such theory by the study of a boundary problem which depends upon an additional parameter ν, in such a way that it reduces to the given (irregular) problem for some specific value of ν, while it remains regular for adjacent values. It is shown that in a specifiable sense the expansions associated with irregular cases of certain categories may be regarded as summable to the given function. (Received August 2, 1940.)

466. W. G. Madow: *Contributions to the theory of the representative method of sampling.*

The theory of representative sampling may be regarded as a dual process, the first consisting in sampling different random variables and the second in repeating several times the experiments associated with each of the different random variables. It follows that while the theory of sampling from finite populations without replacement may be required for the first process, the second leads directly into the theory of sampling from infinite populations. There is, however, one difference. The usual theory is concerned with the evaluation of fiducial or confidence limits for, say, the mean of a sample of N when n, $(N \geqq n)$, of the values are known. It is thus possible to use the usual theories of estimation in obtaining estimates of the parameters and to allow the effects of the subsampling process to show themselves in the different values of the fiducial limits. It is shown that the limits obtained are almost identical with those obtained by the theory of sampling from a finite population. Distributions of the statistics used in these limits are derived. Besides these results, the theory is extended to that of sampling vectors and the conditions are stated under which the "best" allocation of the number in a sample among several strata is proportional to the kth roots of the generalized variance of a random vector having k components. (Received August 2, 1940.)

467. Jerzy Neyman: *Conception of equivalence in the limit of statistical tests and its application to certain forms of χ^2 and to the tests based on the λ principle.*

Denote by T_1 and T_2 two different tests of the same hypothesis H and by N the number of observations to be used to test H. Definition: if the probability of T_1 and T_2 contradicting themselves tends to zero as N is increased, then T_1 and T_2 are called equivalent in the limit. Consider an experiment which will produce one of the results $E_1, E_2, \cdots, E_s$, the probability of E_i being p_i. This experiment is to be repeated n times and n_i will denote the number of occurrences of E_i. Denote by H the hypothesis that $p_i = f_i(\theta_1, \theta_2, \cdots, \theta_h)$, where f_i is a function of h parameters θ_j, with continuous derivatives of second order such that $\partial(f_1, f_2, \cdots, f_h)/\partial(\theta_1, \theta_2, \cdots, \theta_h) \neq 0$. Suppose that H is tested against a set of alternatives ascribing to the p's any non-negative values, such that $\sum_i p_i = 1$. Under those conditions each of the following χ^2 tests is equivalent in the limit to the test based on the λ principle: (1) test T_1 consists in the rule of rejecting H whenever $\sum_i (n_i - np_i')^2/np_i' > \chi^2_\alpha$, where p_i' denotes the maximum likelihood estimate of p_i; (2) test T_2 consists in rejecting H whenever

$\sum_i (n_i - n\bar{p}_i)^2/n\bar{p}_i > \chi^2_\alpha$, where $\bar{p}_i$ denotes the value of p_i minimizing $\sum_i (n_i - np_i)^2/np_i$; (3) test T_3 consists in rejecting H whenever $\sum_i (n_i - np_i'')^2/n_i > \chi^2_\alpha$, where p_i'' denotes the value of p_i minimizing $\sum_i (n_i - np_i)^2/n_i$. (Received September 11, 1940.)

468. Isaac Opatowski: *On the motion of an electric particle.*

The motion of an electrically charged point mass under the action of a magnetic force $\mathfrak{H} = -\text{grad } U(P)$ and an electric force $\mathfrak{E} = -\text{grad } V(P)$ is studied in the case in which U and V are harmonic functions satisfying the conditions: (1) grad $W(P) = \mathfrak{u} \times \mathfrak{H}$, $C = \mathfrak{u} \cdot \mathfrak{E}$ where W is a certain function (generalization of the ordinary stream function), C is a scalar constant, $\mathfrak{u} = \mathfrak{a} + \mathfrak{b} \times \mathfrak{r}$, where $\mathfrak{a}$, $\mathfrak{b}$ are two constant vectors and $\mathfrak{r}$ is the radius vector. The most important fields of mathematical physics, that is, general plane fields, all fields symmetric about an axis of revolution and many others are included in (1). The integral of momentum is explicitly given in terms of W and C. Regions of space are determined within which, for a given energy and a given constant of momentum integral, no motion is possible (generalization of Stoermer's "forbidden regions"). Equations of certain trajectories are explicitly given. The paper deals with both constant and relativistic mass. (Received August 2, 1940.)

469. Gordon Pall: *On the arithmetic of ternary quadratic forms.*

The fundamentals of the arithmetic of ternary quadratics is presented with modifications of the usual invariants, leading to very considerable simplifications. Among the topics treated are: order invariants, form-residues, generic characters, five equivalent definitions of genus, the number of existing genera, simultaneous representation, simultaneous characters, the numbers represented by a genus, zero and universal forms, rational equivalence, reduction formulas for the representations of pn or p^2n, and so on. (Received August 5, 1940.)

470. Everett Pitcher: *The cap heights of a sum function on a product space.*

This paper contains a solution under some restrictions of a problem in critical point theory arising in the work of M. Morse and C. Tompkins (see Annals of Mathematics, (2), vol. 38 (1937), pp. 386–449 and vol. 40 (1939), pp. 443–472, and forthcoming papers) on the problem of Plateau. M and N denote metric spaces and $f(x)$ and $g(y)$ positive functions defined upon them. It is assumed that (i) the sets $f \leqq c < \infty$ are compact, and (ii) the cap heights of f form an increasing sequence, finite or with $+\infty$ as limit. It is assumed that g satisfies (i) and (ii) also. On $M \times N$, $h[x \times y] = f(x) + g(y)$ and satisfies (i) and (ii). Vietoris topology with coefficients from a field is used. An arbitrary maximal group of k-caps of h at height e is shown to be isomorphic to a direct sum of homology groups $H^k(c, d)$. The summation extends over all pairs c, d of cap heights of f and g respectively with sum e. The group $H^k(c, d)$ is the homology group of classes of k-cycles on the relative product space $[f \leqq c] \times [g \leqq d] \bmod [f \leqq c] \times [g < d] \cup [f < c] \times [g \leqq d]$. Some consideration is given to homology groups of a product space. (Received August 6, 1940.)

471. Hillel Poritsky and H. D. Snively: *Doubly periodic networks.*

A solution is obtained for the doubly infinite number of linear equations of a doubly periodic network. Expressions are obtained for the self and mutual impedances as double integrals which are then reduced to elliptic integrals (of all three kinds).

These impedances are also evaluated by direct numerical methods of successive approximation. (Received August 21, 1940.)

472. J. F. Randolph: *Some properties of sets of the Cantor type.*

Given $a_1 \geqq a_2 \geqq \cdots > 0$ where $\sum_n a_n = 1$, let S consist of the numbers x for which there exists a sequence $\{\epsilon_n\}$, where each ϵ_n is 0 or 1, such that $x = \sum_n \epsilon_n a_n$. (If $a_n = 2/3^n$ then S is the Cantor set.) Theorem: $a_n \geqq 2\sum_{i=n+1}^{\infty} a_i$ is a necessary and sufficient condition that corresponding to $0 \leqq z \leqq 2$ there exists an $x \in S$ and $y \in S$ such that $x + y = z$. Also some results are given on the Carathéodory and Gillespie linear measures of cartesian products of sets of the Cantor type. (Received August 6, 1940.)

473. W. T. Reid: *A new class of self-adjoint boundary value problems.*

This paper treats a differential system consisting of the vector differential equation $\mathcal{L}[y] \equiv y' - A(x)y = \lambda B(x)y$ and two-point boundary conditions $s[y] \equiv My(a) + Ny(b) = 0$ which is self-adjoint under the nonsingular transformation $z = T(x)y$ and satisfies the further conditions: (1) the matrix $S(x) = \tilde{T}(x)B(x)$ is symmetric on ab; (2) $H[y] = \int_a^b y\tilde{T}\mathcal{L}[y]dx > 0$ for all vectors y satisfying $s[y] = 0$, $B(x)y(x) \not\equiv 0$, and for which there exists a corresponding vector $g(x)$ such that $\mathcal{L}[y] = Bg$ on ab; (3) $y \equiv 0$ is the only vector satisfying on ab the conditions $\mathcal{L}[y] = 0$, $By \equiv 0$, $s[y] = 0$. The chief distinction between these conditions and those defining a definitely self-adjoint boundary problem (Bliss, Transactions of this Society, vol. 44 (1938), pp. 413–428) is that the above hypothesis (2) on $H[y]$ replaces a corresponding definiteness assumption on the matrix $S(x)$. Such systems are shown to possess fundamental properties similar to those previously established for definitely self-adjoint problems. In particular, certain important types of boundary problems associated with the calculus of variations which are not definitely self-adjoint do belong to this new class of problems. (Received August 6, 1940.)

474. I. J. Schoenberg: *On positive definite functions on spheres.*

The positive definite functions on m-dimensional spheres and on spheres in Hilbert space are explicitly determined. In ordinary terminology the results are as follows: I. Let $f(x)$ be real and continuous in the interval $-1 \leqq x \leqq 1$. Let $A = \|a_{ik}\|$ denote an arbitrary n-rowed symmetric matrix of a positive quadratic form whose rank does not exceed the fixed number $m+1$ ($m \geqq 1$). Moreover $a_{11} = a_{22} = \cdots = a_{nn} = 1$. The quadratic form corresponding to the transformed matrix $f(A) = \|f(a_{ik})\|$ is also always positive if and only if $f(x)$ admits throughout $[-1, 1]$ an expansion in ultraspherical polynomials $f(x) = \sum_{\nu=0}^{\infty} p_\nu P_\nu^{((m-1)/2)}(x)$ with nonnegative coefficients $p_\nu \geqq 0$. II. The assumptions on $f(x)$ and the matrix A are as above except that there is now no restriction whatever on the rank of A. The quadratic form of $f(A) = \|f(a_{ik})\|$ is also always positive if and only if $f(x) = p_0 + p_1 x + p_2 x^2 + \cdots$, $(-1 \leqq x \leqq 1;\ p_\nu \geqq 0)$. (See Pólya and Szegö, vol. 2, p. 107, where the sufficiency of the power series expansion for $f(x)$ is stated as Problem 37.) The second theorem is derived from the first by letting $m \to \infty$. This passage to the limit is carried through by means of the following new estimate: $|P_n^{(\lambda)}(\cos\theta)| \leqq P_n^{(\lambda)}(1)\{((1+\cos^2\theta)/2)^{n/2} + (4/n)\cdot(1+\cos^2\theta)/\sin(2\theta)\}$ if $0 < \theta < \pi/2$, $\lambda \geqq 5$, and $n = 1, 2, 3, \cdots$. (Received August 5, 1940.)

475. I. M. Sheffer: *On the singularities of functions under certain linear transformations.*

The linear transformations under consideration are all of the form $L[y(x)]$

$=\sum_0^n L_n(x)y^{(n)}(x)$. When $L[y(x)]=F(x)$, the relation between the singularities of $F(x)$ and those of $y(x)$ is examined. The theorems obtained are analogous to the well known theorems of Hadamard and Hurwitz on the singularities of power series compounded from given power series; indeed, these latter theorems are but particular instances of the results of the present work. (Received August 6, 1940.)

476. D. C. Spencer: *On an inequality of Grunsky.*

Let W be a Riemann domain which, for simplicity, is assumed to be the map of $|z|<1$ by a function $f(z)$ regular in $|z|\leqq 1$, and let $n(w)$ be the number of times the point w is covered by W. Suppose that $n(w)\leqq n(0)$. Then Grunsky (Dissertation, Schriften des mathematischen Seminars der Universität Berlin, vol. 1 (1932), pp. 113–118) proves the following inequality (a case of the theorem of the geometric and arithmetic means in which the weight function is not necessarily of constant sign): $\exp\{(1/2\pi n(0))\int_B \lg R^2 d\Phi\}\leqq(1/2\pi n(0))\int_B R^2 d\Phi$, where B is the boundary of W. In this note a new and simple proof of the inequality is given for more general domains W which are only mean $n(0)$-valent (that is, under conditions less restrictive than $n(w)\leqq n(0)$). The inequality is applied to obtain a more precise version of a theorem of Bermant (Comptes Rendus de l'Académie des Sciences, Paris, vol. 207 (1938), pp. 31–33). (Received August 6, 1940.)

477. Gabor Szegö: *Power series with multiply monotonic sequences of coefficients.*

A sequence $\{a_0, a_1, \cdots\}$ is called monotonic of order k if the differences $\Delta^{(\nu)}a_n$ defined by $\Delta^{(0)}a_n=a_n$, $\Delta^{(1)}a_n=a_n-a_{n+1}$, $\Delta^{(\nu)}=\Delta^{(1)}\Delta^{(\nu-1)}$ are non-negative for $0\leqq\nu\leqq k$, $n\geqq 0$. Let $a_n\not\equiv 0$. Fejér proved (Transactions of this Society, vol. 39 (1936), p. 57) that $\sum a_n z^n$ is regular and univalent for $|z|<1$ provided $\{a_n\}$ is monotonic of order $k=4$. He also showed that this is not true for $k=1$. In the present paper it is proved that the assertion in question remains true for $k=3$, but not for $k=2$. The latter fact has been established by a different method by Mr. S. Szidon (Acta Szeged, 1940). (Received September 11, 1940.)

478. W. J. Trjitzinsky: *Properties of growth for solutions of differential equations of dynamical type.*

In the background of this work are, in part, certain memoirs of A. Liapounoff, P. Bohl, E. Cotton, and O. Perron. The developments for linear systems are based on characteristic numbers and product integration. The nonlinear systems are investigated, in part, with the aid of the theory of linear systems. In the study of the nonlinear problem, use is made of characteristic numbers and of successive approximations. The latter method is associated with some of the present author's earlier work in the field of differential equations. (Received August 5, 1940.)

479. Abraham Wald: *Asymptotically shortest confidence intervals.*

Let $f(x, \theta)$ be the probability density function of a variate x involving an unknown parameter θ. Let $x_1, x_2, \cdots, x_n$ be n independent observations on x and let $C_n(\theta)$ be a positive function of θ such that the probability that $F(x, \theta)\equiv\left|n^{-1/2}\partial\sum_{\alpha=1}^{n}\log f(x_\alpha, \theta)/\partial\theta\right|\leqq C_n(\theta)$ is a constant β when θ is the true value of the parameter. Denote by $\underline{\theta}(x_1, \cdots, x_n)$ the root in θ of the equation $F(x, \theta)=C_n(\theta)$ and by $\bar{\theta}(x_1, \cdots, x_n)$ the root of $F(x, \theta)=-C_n(\theta)$. Under some weak assumptions on $f(x, \theta)$ the limiting

interval $\delta_n(x_1, \cdots, x_n) = [\underline{\theta}(x_1, \cdots, x_n), \overline{\theta}(x_1, \cdots, x_n)]$ for $n \to \infty$ is a shortest unbiased confidence interval (for the definition of a shortest unbiased confidence interval, see J. Neyman, Philosophical Transactions of the Royal Society of London, vol. 236 (1937), pp. 333–380) of θ corresponding to the confidence coefficient β. This confidence interval is identical with that given by S. S. Wilks (Annals of Mathematical Statistics, vol. 9 (1938)). Wilks has shown that $\delta_n(x_1, \cdots, x_n)$ is asymptotically shortest in the average compared with all confidence intervals computed on the basis of statistics belonging to a certain class C. In the present paper it has been proved that the confidence interval in question is asymptotically shortest compared with any arbitrary unbiased confidence interval, without any restriction to a certain class of functions. (Received August 5, 1940.)

480. Norbert Wiener and Aurel Wintner: *Discrete chaos.*

The ordinary method of introducing the Lebesque integration needed in statistical mechanics in terms of a probability density referred to a product space works in systems of a finite number of degrees of freedom, but breaks down in the case of an infinity of degrees of freedom. An alternative method is developed in this latter case, depending on an infinity of probability densities of finite sets of particles. (Received September 11, 1940.)

481. S. S. Wilks: *On the problem of two samples from normal populations with unequal variances.*

Suppose O_{n_1} and O_{n_2} are samples of n_1 and n_2 elements from normal populations Π_1 and Π_2 respectively. Let a_1, σ_1^2 and a_2, σ_2^2 be the means and variances of Π_1 and Π_2 and let O_{n_1} and O_{n_2} have means $\bar{x}_1$ and $\bar{x}_2$ and variances s_1^2 and s_2^2 (unbiased estimates of σ_1^2, σ_2^2) respectively. It is shown that there exists no function (Borel measurable) of $\bar{x}_1$, $\bar{x}_2$, s_1^2, s_2^2, $a_1 - a_2$ independent of σ_1^2 and σ_2^2, having its probability law independent of the four population parameters. It is therefore impossible to obtain exact confidence limits for $a_1 - a_2$ corresponding to a given confidence coefficient. Functions of the four parameters and four statistics are devised from which one can set up confidence limits for $a_1 - a_2$ with associated confidence coefficient inequalities. (Received August 2, 1940.)

482. W. L. G. Williams: *The application of hyperbolic complex numbers to the geometry of the triangle.*

A complex number $a + bj$, in which a and b are real numbers and $j^2 = 1$, is called a hyperbolic complex number. If such a number is designated by z and is represented as a two-dimensional vector, the vector zj is said to be pseudo-perpendicular to z. When the concept pseudo-perpendicular takes the place of perpendicular, new points, related to a triangle, are found, analogous to orthocentre, circumcentre and nine-point centre. These points lie on a straight line with the centroid of the triangle. There exist rectangular hyperbolas analogous to the circumcircle and nine-point circle. There is an analogue of Feuerbach's theorem. (Received August 6, 1940.)

483. Antoni Zygmund: *Power series of the class H^λ on the circle of convergence.*

If the function $f(z) = \sum c_n z^n$ belongs to the class H^λ, where $0 < \lambda < 1$, then the series $\sum c_n e^{in\theta}$ is summable $(C, 1/\lambda - 1)$ for almost every θ. (Received August 19, 1940.)

484. A. T. Brauer: *On a problem of partitions.*

Let $a_1, a_2, \cdots, a_k$ be relatively prime positive integers. The purpose of this paper is to determine bounds $F(a_1, a_2, \cdots, a_k)$ such that the Diophantine equation $a_1x_1+a_2x_2+\cdots+a_kx_k=n$ always has solutions in positive integers $x_1, x_2, \cdots, x_k$ for $n>F(a_1, a_2, \cdots, a_k)$. (Received September 27, 1940.)

485. A. B. Brown: *On the number of independent parameters.*

Let $f(x_1, \cdots, x_n, \alpha_1, \cdots, \alpha_m)$ be a function of class $C^{(q)}$ (q specified in the paper). To form matrix (M), take $\partial f/\partial\alpha_1, \cdots, \partial f/\partial\alpha_m$ as first row. As further trial rows, take the derivatives to $\alpha_1, \alpha_2, \cdots, \alpha_m$ of derivatives of f to the x's, in the following order: $\partial f/\partial x_n, \partial f/\partial x_{n-1}, \cdots, \partial f/\partial x_1, \partial f/\partial x_n^2, \cdots$. If the set of derivatives to $\alpha_1, \cdots, \alpha_m$ increases the rank of the matrix, the derivative, say $\partial z/\partial x_n$, is *accepted*, otherwise *rejected*. The algorithm continues until the accepted derivatives are the only derivatives of z which are not rejected derivatives or derivatives of rejected derivatives. If k is the number of rows of the resulting matrix, then *all functions f are given by varying k of the parameters, keeping the other $n-k$ constant; furthermore, no function is given twice in this way.* Certain "singular points," forming a nowhere dense closed set, must be avoided. It is proved that not all functions can be obtained by varying less than k parameters. Further, given $g(x_1, \cdots, x_n, \beta_1, \cdots, \beta_l)$ where $f(x, \alpha)$ and $g(x, \beta)$ are the same set of functions of (x), but nothing is assumed as to relationship between (α) and (β), the same number k of parameters is needed for $g(x, \beta)$. (Received September 25, 1940.)

486. M. M. Day: *Some more uniformly convex spaces.*

For a sequence of Banach spaces $[B_i]$, $i=1, 2, \cdots$, define $B=\prod^p\{B_i\}$ to be the space of sequences $b=\{b_i\}$ with $b_i\in B_i$ and $\|b\|=(\sum\|b_i\|^p)^{1/p}<\infty$. Complementing an earlier result of this writer (abstract 46-9-396) it is shown that $\prod^p\{B_i\}$ is uniformly convex if $B_i=l^{p_i}$ or L^{p_i} and if $1<m\leqq M<\infty$ exist with $m\leqq p_i\leqq M$ for all i. A conjecture of Boas (this Bulletin, vol. 46 (1940), p. 304) that the spaces $\prod^p\{B_i\}$ with $B_i=B_0$ for all i, as well as the spaces $L^p(B_0)$ (consisting of all Bochner integrable functions f on, say, (0, 1) with values in B_0 and $\|f\|=(\int_0^1\|f(t)\|^p dt)^{1/p}<\infty$), are uniformly convex if B_0 is shown to be true. A sequence of spaces $\{B_i\}$ is said to have a common modulus of convexity if for each ϵ, $0<\epsilon\leqq 2$, there is a $\delta(\epsilon)>0$ such that for each i and each pair of points b_i, $b_i'\in B_i$ with $\|b_i\|=\|b_i'\|=1$ and $\|b_i-b_i'\|>\epsilon$, the relation $\|b_i+b_i'\|<2(1-\delta(\epsilon))$ holds. The results mentioned above are corollaries of the following theorem: $\prod^p\{B_i\}$ is uniformly convex if and only if the B_i have a common modulus of convexity. (Received September 21, 1940.)

487. Aaron Fialkow: *The conformal theory of a hypersurface.*

For a hypersurface V_{n-1} in any Riemann space V_n ($n>2$), the author shows that it is possible to define three quadratic differential forms which remain unchanged under any conformal mapping of V_n, not necessarily on itself. These forms are the *conformal fundamental forms* of the hypersurface V_{n-1}. If $n>3$, the coefficients of the third conformal fundamental form may be expressed in terms of the coefficients of the first and second conformal fundamental forms. If V_{n-1} and $\bar{V}_{n-1}$ are hypersurfaces of V_n and $\bar{V}_n$ respectively and $V_n\leftrightarrow\bar{V}_n$, $V_{n-1}\leftrightarrow\bar{V}_{n-1}$ by a conformal map, then the conformal fundamental forms of V_{n-1} and $\bar{V}_{n-1}$ are equal. Conversely, if V_n and $\bar{V}_n$ are conformally euclidean spaces and the conformal fundamental forms of V_{n-1} and $\bar{V}_{n-1}$ are equal, then a conformal transformation exists so that $V_n\leftrightarrow\bar{V}_n$, V_{n-1}

$\leftrightarrow \bar{V}_{i-1}$. In any V_n, a hypersurface exists whose conformal fundamental forms are any preassigned quadratic differential forms whose coefficients satisfy certain partial differential equations analogous to the classical Gauss-Codazzi equations. The Weyl conformal curvature tensor of V_n plays a role analogous to that of the Riemann curvature tensor in classical differential geometry. The case $n=3$ occupies a special position in this theory. (Received September 23, 1940.)

488. G. E. Forsythe: *On Riesz summability methods of order r, for* $R(r)<0$. Preliminary report.

Let A_r and B_r represent M. Riesz's (discrete *resp.* continuous) summability methods for sequences, as defined by Agnew (Transactions of this Society, vol. 35 (1933), pp. 532–548). It is known that A_r and B_r and Cesàro C_r are equivalent for $-1<r\leqq 1$, while A_r and B_r are not equivalent for the real part of r, $R(r)<-1$, nor for certain values of $r>1$. By aid of Theorem 5.1 (op. cit.) the present paper obtains a criterion for the equivalence of A_r and B_r for $R(r)<0$ in terms of the reciprocal of $\phi_r(x)=\sum_1^\infty n^r x^n$, and applies it to prove that A_{-1+ih} and B_{-1+ih} $(-\infty<h<\infty)$ are equivalent if and only if $h=0$. (Received September 25, 1940.)

489. Saul Gorn: *Homomorphisms and modular functionals.*

Using CA to represent the complements of elements of A, an ideal in a complemented modular L is called a C-ideal if $C\mathfrak{A}$ is an ideal; $C\mathfrak{A}$ is a π-ideal if $\mathfrak{A}$ is a σ-ideal and dually. If $\mathfrak{A}=CC\mathfrak{A}$, $\mathfrak{A}$ is called neutral (identical with Garrett Birkhoff's definition). An $\mathfrak{A}$ fulfilling both conditions is called C-neutral. The general homomorphism theorem for complemented modular lattices is given by the C-neutral ideals. Letting $\mathfrak{A}'$ be the elements orthogonal to all elements of $\mathfrak{A}$, a C-neutral ideal is normal if and only if it is normal in Stone's sense: $\mathfrak{A}''=\mathfrak{A}$. The homomorphism $L\rightarrow L/\mathfrak{A}$ takes normal ideals containing or equal to $\mathfrak{A}$ into normal ideals; conversely if $\mathfrak{A}$ is normal. L is called ideally irreducible if it contains no non-trivial C-neutral normal ideal. $L/\mathfrak{A}$ is ideally irreducible if and only if $\mathfrak{A}$ is a maximal (or prime for π-ideals) C-neutral normal ideal. Using the Wilcox-Smiley continuity conditions, a quasi-norm is uniquely determined by its generating normal C-neutral $\mathfrak{A}$ if and only if $L/\mathfrak{A}$ is ideally irreducible (that is, $\mathfrak{A}$ maximal if non-trivial). This generalizes results mentioned in abstract 45-1-16. (Received September 12, 1940.)

490. N. A. Hall: *The solution of the quadrinomial equation in infinite series by the method of iteration.*

The roots of the quadrinomial equation $z^{m+n+k}-z^{m+n}+bz^m+a=0$ may be expressed as double power series in the two complex variables a and b. These solutions have been previously developed by use of the Lagrange expansion and also by use of differential resolvents. In this paper, these results are reproduced and extended by the direct method of solution of the equation by iteration as previously applied by the author to the trinomial equation (this Bulletin, vol. 44 (1938), p. 337). The solutions are exhibited as generalized hypergeometric series in the two variables a and b. These series give the set of $m+n+k$ roots covering the quadrant of absolute values of a and b except for the small region covered by the branch points of the roots. The relation between these singularities and the domains of convergence for the double series is shown. (Received September 27, 1940.)

491. Einar Hille: *A class of differential operators of infinite order.*

The author studies the differential operator $G(\delta_x)$ where $G(w)$ is an entire function

and $\delta_z = z^2 - d^2/dz^2$ is the operator of the Hermite-Weber equation. Necessary and sufficient conditions are found in order that $G(\delta_z)$ shall apply to various classes of analytic functions, in particular entire functions. The conditions differ in some respects fundamentally from corresponding conditions for the operator $G(d/dz)$. Relations are found between the orders and types of $G(w)$, $f(z)$, and $G(\delta_z)\cdot f(z)$. The equation $G(\delta_z)\cdot W = F(z)$ is studied with the aid of summable Fourier-Hermite series. If $F(z) \sim \sum F_n h_n(z)$, then formally $W(z) \sim \sum W_n h_n(z)$, where $G(2n+1)W_n = F_n$. Sufficient conditions are found under which this formal procedure leads to an actual solution and a uniqueness theorem is proved. Some examples are discussed which show that if the operator δ_z is replaced by other second order differential operators D_z then the applicability of $G(D_z)$ to entire functions may be governed by completely different conditions. (Received August 30, 1940.)

492. Loo-Keng Hua: *Some "Anzahl" theorems in the theory of groups of prime-power order.*

A group G of order p^n (p being a prime) is said to be of rank δ if the maximum of the order of elements of G is equal to $p^{n-\delta}$. The author selects the following theorem to be announced: If G is a group of order p^n ($p \geqq 3$, $u \geqq 2\alpha$) of rank α, then (i) G contains one and only one subgroup of order p^m ($\alpha \leqq m \leqq n$) of rank α; (ii) G contains p^{α} cyclic subgroups of order p^m ($\alpha < m < n-\alpha+1$) and (iii) G contains $p^{m+\alpha}$ ($\alpha \leqq m \leqq n-\alpha$) elements which satisfy $x^{p^m} = 1$. The second and the third statement solve the "Anzahl" theorems of Miller's and Kalakoff's type. (Received August 31, 1940.)

493. Loo-Keng Hua: *The lattice points in a circle.*

Let $R(x)$ denote the number of lattice points in the circle $m^2+n^2 = x$. The object of the paper is to prove that $R(x) = \pi x + O(x^{\alpha+\epsilon})$, $\alpha = 13/40$, which is better than Titchmarsh's result (Proceedings of the London Mathematical Society, (2), vol. 38 (1933), pp. 96–115) with an error term $O(x^{15/46+\epsilon})$. Notice that Vinogradow's proof of the error term $O(x^{17/53+\epsilon})$ is incorrect (Bulletin de l'Académie des Sciences de l'URSS, vol. 7 (1932), pp. 313–336). (Received August 31, 1940.)

494. W. H. Ingram: *A generalization of Erhard Schmidt's solution of the nonhomogeneous integral equation.* Preliminary report.

It is first proved that the Hilbert transform of an arbitrary step-wise continuous vector matrix has, in terms of the characteristic solutions of the equation $\phi_r(x) = \lambda_r \cdot \int_0^1 \mathfrak{K}(x, \alpha)\phi_r(\alpha)d\alpha$, an absolutely and uniformly convergent development which represents the transform in the sense of least squares when the scalar coefficients are computed by means of the associated characteristic functions ψ_r. As in abstract 46-5-282, $\mathfrak{K}$ is a non-symmetric square matrix, and ϕ_r and ψ_r, respectively, column and line matrices which can be biorthonormalized. In this sense of equality and on the basis of this expansion theorem, the generalization of Schmidt's result previously announced is independently derived. (Received August 27, 1940.)

495. L. H. Loomis: *The decomposition of meromorphic functions into rational functions of univalent functions.*

The general question discussed is this: given $f(z)$ meromorphic in $|z| < 1$, when does there exist a decomposition $f(z) \equiv f_2(f_1(z))$ where $\zeta = f_1(z)$ is univalent in $|z| < 1$, and $f_2(\zeta)$ is a rational function? A simple but typical result is that if $f(z)$ is meromorphic in $|z| \leqq 1$, then the decomposition is possible. Necessary and sufficient conditions

are derived for the general case. Stronger theorems are proved for situations in which $\zeta = f_1(x)$ is the mapping function of a simply-connected plane region which is not of the most general type, for instance, a Jordan region. Analogous theorems are developed for $f(z)$ regular and bounded in $|z| < 1$ and $f_2(\zeta)$ restricted to special types of rational functions, say polynomials. In proof, the author considers the possibility of imbedding the Riemann surface which is the map of $|z| < 1$ under $w = f(z)$ in the Riemann surface of the inverse of a rational function. (Received September 23, 1940.)

496. P. T. Maker: *A topological characterization of monotone functions.* Preliminary report.

Let $f(s)$, a function from S to X, both topological spaces, be in $M(S, X)$ if the complete inverse image of every connected set in X is connected. It is shown that $M(R_1, R_1)$ is the class of monotone functions, and that for other spaces M includes the functions "monotone" by previous generalizations. Conditions on S and X, in order that $f(s)$ have certain continuity properties, or that M be complete or compact, are determined. (Received September 28, 1940.)

497. E. R. Ott: *A locus determined by an algebraic correspondence.*

Let P and Q be two variable vertices of a hexagon inscribed in a conic and let the other four vertices be fixed. If P and Q have parameters $t' = F(t)$ and $t' = G(t)$, where $F(t)$ and $G(t)$ are functions which establish an (m, n) algebraic correspondence between P and Q, then as t varies the Pascal line of the hexagon envelopes a curve of class $(m+n)$ and of order $2(m+n-1)$. Each of the two fixed sides of the hexagon which are adjacent to the variable side is a multiple tangent of the locus. All of the Pluecker characteristics of the curve are obtained and are independent of the order of the vertices of the hexagon. (Received September 14, 1940.)

498. Francisco Perez: *A generalization of the theory of invariant factors and similar matrices.*

E. H. Moore's general identity matrix $\delta_{\mathfrak{PP}}$ (E. H. Moore, *General Analysis*, Part I) for a right linear space $\mathfrak{M}$ of vectors on a finite range $\mathfrak{P}$ to $\mathfrak{A}$ is shown to exist even when $\mathfrak{A}$ is a number system of *type B*, provided $\mathfrak{M}$ has the property of being *perfect*. In a perfect space $\mathfrak{M}$ of rank r, to every matrix ϕ of type $\mathfrak{M}\overline{\mathfrak{M}}$ and of rank r corresponds uniquely a matrix ϕ^{-1} of type $\mathfrak{M}\overline{\mathfrak{M}}$ such that $S\phi^{-1}\phi = S\phi\phi^{-1} = \delta_{\mathfrak{PP}}$. When multiplication is commutative, it is shown (without using determinants) that every matrix κ of type $\mathfrak{M}\overline{\mathfrak{M}}$ determines uniquely a set of polynomials $P_i(z)$ $(i = 1, 2, \cdots, t)$ called the *invariant factors of* κ *in* $\mathfrak{M}$. The sum of the degrees of these polynomials is equal to r, and P_{i+1} divides P_i. It is shown that $P_1(\kappa) = O(\mathfrak{P}\mathfrak{P})$, where $\kappa^0 \equiv O(\mathfrak{P}\mathfrak{P})$. Two matrices κ_1 and κ_2 of type $\mathfrak{M}\overline{\mathfrak{M}}$ are said to be *similar in* $\mathfrak{M}$ if there exists a matrix ϕ of type $\mathfrak{M}\overline{\mathfrak{M}}$ and of rank r such that $\kappa_2 = SS\phi^{-1}\kappa_1\phi$. Two matrices of type $\mathfrak{M}\overline{\mathfrak{M}}$ are similar in $\mathfrak{M}$ if and only if they have the same invariant factors in $\mathfrak{M}$. (Received September 14, 1940.)

499. L. B. Robinson: *A functional equation with a singular line.*

The equation (I) $u'(x) = a(x)u(x^2)$ can be transformed into (II) $w'(z) = e^z a(e^z)w(2z)$ which in general has a solution with $z = 0$ as singular point. The exceptional case is given by the vanishing of a determinant. In general the circumference of the unit circle is a singular line of $u(x)$. The necessary and sufficient condition for the failure of this rule has been found. (Received September 14, 1940.)

500. L. B. Robinson: *A functional equation with negative exponent.*

Consider the system $u_x'(x)=(\lambda/(1+x^2))[P_n(x)/Q_n(x)]u(x^{-2})$, $u_x'(x^{-1})=(-\lambda/(1+x^2))[P_n(x^{-1})/Q_n(x^{-1})]u(x^2)$. The solution of this system is $u(x)\equiv u_0\{1+r\lambda\phi_1(x)+\lambda^2\phi_2(x)+r\lambda^3\phi_3(x)+\cdots\}$, $u(x^{-1})\equiv u_0\{r+\lambda\psi_1(x)+r\lambda^2\psi_2(x)+\lambda^3\psi_3(x)+\cdots\}$ where $\phi_s(x)\equiv\int_0^x(1/(1+\xi^2))[P_n(\xi)/Q_n(\xi)]\psi_{s-1}(\xi^{-2})d\xi$, $\psi_s(x)\equiv\int_0^x(-1/(1+\xi^2))[P_n(\xi^{-1})/Q_n(\xi^{-1})]\cdot\phi_{s-1}(x^2)d\xi$. If the value of r is rightly determined, the two series $u(x)$ and $u(x^{-1})$ converge and are consistent. (Received September 14, 1940.)

501. Jenny E. Rosenthal: *Generating functions and properties of certain orthogonal polynomials.* II.

The nonhomogeneous differential equation satisfied by generating functions for orthogonal polynomials reduces in the simplest possible case to an algebraic equation. Its solution is the generating function of a certain type of Szegö-Bernstein polynomials. A method is given for obtaining the weight factor from the generating function and for finding some additional properties of these polynomials. The second order differential equation satisfied by the polynomials is derived and is shown to reduce in a special case to an equation which was obtained by Shohat. (Received September 27, 1940.)

502. Peter Scherk: *On real closed curves of the order $n+1$ in the projective n-space.* Preliminary report.

The author discusses closed curves K^{n+1} in the real projective n-space which have osculating spaces of all the dimensions less than n and which have the real order $n+1$; that is, the maximum number of points of intersection of a K^{n+1} with an $(n-1)$-space shall be $n+1$. These curves are a generalization of the algebraic curves of order $n+1$ with one real branch. Analogous to the algebraic case, singular points can be defined and provided with multiplicities. The sum of the multiplicities of the singular points of a K^{n+1} is less than or equal to $n+1$ and congruent to $n+1$ (mod 2). The types for which this sum equals $n+1$ are characterized. Furthermore, the pairs of curves in the projective n-space are described which have no more than $n+1$ points with any $(n-1)$-space in common. These pairs generalize the algebraic curves of order $n+1$ with two real branches. Some special qualities of the K^{n+1} are indicated. The main tool of the discussion is the continuous transformation of the K^{n+1} into itself which correlates to each point of the K^{n+1} the point at which its osculating $(n-1)$-space intersects the K^{n+1} again. (Received September 27, 1940.)

503. W. T. Scott and H. S. Wall: *A geometrical method in the theory of continued fractions.*

In this paper the authors consider continued fractions of the form $1/1+a_2/1+a_3/1+a_4/1+\cdots$, where the a_n's are complex numbers. By regarding the continued fraction as a succession of linear fractional transformations they determine conditions under which the nth approximant A_n/B_n of the continued fraction will lie in a region V of the complex plane whenever the a_n's lie in a region U. Under these conditions any convergent continued fraction whose elements lie in U must have a value in V. As a supplement to their recent "parabola" theorem (Transactions of this Society, vol. 47 (1940), p. 166) they show that if the a_n's lie in the parabola $|z|-R(z)\leqq\frac{1}{2}$ every approximant A_n/B_n lies in the circle $|z|^2-2R(z)\leqq 0$, and that this is the "best" circular region having the above property. (Received August 19, 1940.)

504. W. T. Scott and H. S. Wall: *Linear manifolds of Hausdorff means.*

A set M of moment sequences is a manifold if when $\{c_n\}$, $\{d_n\}$ are in M, then $\{c_n+d_n\}$ and $\{rc_n\}$ (r real) are in M. A manifold M is regular if when $\{c_n\}\in M$, $c_0\neq 0$ then $\{c_n/c_0\}$ is a regular Hausdorff sequence. The authors determine conditions under which a sequence $\{\beta_n(u)\}$ of real functions on the interval $(0, 1)$ constitutes a basis for a manifold $M(\beta)$, in the sense that M is made up of the sequences $c_n=\int_0^1\beta_n(u)\cdot d\phi(u)$, $(n=0, 1, 2, \cdots)$, for $\phi(u)\in BV[0,1]$. They also determine conditions upon the basis such that $M(\beta)$ shall be regular, and conditions such that every regular Hausdorff mean determined by the sequences of $M(\beta)$ shall include a given Hausdorff mean. The general theory is applied to a number of special examples. Manifolds exist which include any given Hausdorff mean, and, in particular, which include (C, α) but not $(C, \alpha+\epsilon)$, $(\alpha>0, \epsilon>0)$. For example, if $\beta_n(u)=(u+1)/(u+n+1)$, then $M(\beta)\supset(C, 1)$ but not $(C, 1+\epsilon)$, $(\epsilon>0)$. (Received August 19, 1940.)

505. I. E. Segal: *The space of Besicovitch almost periodic functions.*

Let B^p $(1\leqq p<\infty)$ be the space of all Besicovitch almost periodic functions of order p on the infinite interval $(-\infty, \infty)$ with the usual norm. It is shown that there is a bicompact space S and a completely additive measure μ over the field of Borel subsets of S such that B^p is isometric with the space $L_p(S, \mu)$ of all complex-valued functions which are measurable and whose pth power is summable with respect to μ; the norm is the usual one. This result follows readily from a recent theorem of Gelfand on the representation of the class of Bohr almost periodic functions. It is also connected with some recent work of Bochner and of Wecken. A similar result holds for the spaces of almost periodic functions on any locally bicompact group. (Received September 28, 1940.)

506. William Wernick: *Distributive properties of set operators.*

A point set operator a which maps a set A of a given space S uniquely into a set $aA\in S$ may have certain distributive properties defined by expressions of the form $a(Af_1B)\cdot R:aA\cdot f_2\cdot aB$; where f_1, f_2 are set operators $+$ or $\cdot$; and where R is a set relation, either $=$, $\subset$, or $\supset$. There are twelve such properties, but *monotonicity* and *inverse monotonicity* are included to obtain a list of fourteen "distributive" properties. These fourteen properties (α_i) and their negations $(\bar{\alpha}_i)$ form a collection of 28 properties which may be assumed or deduced for set operators. Their interrelations are investigated. The distributive character of a is determined with respect to α_i if it is known that a has property α_i $(a:\alpha_i)$; or that $a:\bar{\alpha}_i$. An operator a is "completely determined" if its distributive character is determined with respect to every α_i. In view of the interrelations among the α_i, there are relatively few "completely determined" operators. A list of 25 is given, with examples. (Received September 13, 1940.)

507. William Wernick: *Functional dependence in the calculus of propositions.*

In the two-valued calculus of propositions a function F of n variables is completely determined by 2^n constants, independently 0, or 1 (call their sum S). F is independent of a particular variable x_i if no change in F results from a change in x_i alone. A function A_i is defined whose vanishing is a necessary and sufficient (NS) condition that F be independent of x_i. In terms of the A_i a function $A(F)$ is defined whose nonvanishing

is a NS condition that F be really a function of all its arguments, that is, not independent of any of them. Another function B_i is defined, and it is shown that $S=2B_i$ is another NS condition that F be independent of x_i. From this are obtained very simply the following two theorems: "A sufficient condition that F be really a function of all its arguments is that S be odd." "A necessary condition that F be independent of some or all of its arguments is that S be even." (Received September 20, 1940.)

508. Hassler Whitney: *The mappings of a 3-complex into a space with vanishing fundamental group.*

Extending methods used formerly for classifying the mappings of a 3-complex into a 2-sphere S^2 (see this Bulletin, vol. 42 (1936), p. 338), S^2 is now replaced by any space S without fundamental group. The 2-dimensional homotopy group $\pi^2(S)$ is a direct sum of cyclic groups, with generators $\rho_1, \cdots, \rho_n$. There is a natural function $\theta(\rho)\in\pi^3(S)$ and a multiplication $\rho\times\rho\in\pi^3(S)$; $2\theta(\rho)=\rho\times\rho$. If $a\rho=0$, then $2a\theta(\rho)=0$. If f is a mapping of the 2-dimensional part K^2 of K into S, a 2-π^2-chain X^2 is defined (compare Duke Mathematical Journal, vol. 3 (1937), pp. 51–55); f may be extended through K^3 and K^4 if and only if $\delta X^2=0$ and $X^2\cup^* X^2\sim 0$. Here, if $\sigma^r\cup\sigma^s=\sigma^{r+s}$, then $\sum\alpha_i\rho_i\sigma^r \cup^*\sum\beta_i\rho_i\sigma^s=[\sum\alpha_i\theta(\rho_i)+\sum_{i<j}\alpha_i\beta_j(\rho_i\times\rho_j)]\sigma^{r+s}$ if no ρ_i are of finite order. Two mappings f and f' are homotopic if and only if their difference is $\sim 2(Y^1\cup^* X^2)$ for some 1-π^2-cocycle Y^1. (The factor 2 was omitted in the abstract cited above.) (Received September 28, 1940.)

NEW PUBLICATIONS

BELL, E. T. The Development of Mathematics. New York and London, McGraw-Hill, 1940. 13+583 pp.

BRITISH ASSOCIATION FOR THE ADVANCEMENT OF SCIENCE. Mathematical Tables. Vol. 8. Number-Divisor Tables. Cambridge, University Press; New York, Macmillan, 1940. 10+100 pp.

BURINGTON, R. S. Handbook of Mathematical Tables and Formulas. 2d edition. Sandusky, Ohio, Handbook Publishers, 1940. 275 pp.

ELLIOTT, W. W., and MILES, E. R. C. College Mathematics. A First Course. New York, Prentice-Hall, 1940. 13+396 pp.

GILL, T. B. See LANGER, C. H.

GÖDEL, K. The Consistency of the Axiom of Choice and of the Generalized Continuum-Hypothesis with the Axioms of Set Theory. (Annals of Mathematics Studies, no. 3.) Princeton, University Press; London, Humphrey Milford and Oxford University Press, 1940. 66 pp.

LANGER, C. H., and GILL, T. B. Mathematics of Accounting and Finance. Part 1. 26+495+114 pp. Part 2. 19+542+170 pp. Chicago, Walton, 1940.

LIEBER, H. G. See LIEBER, L. R.

LIEBER, L. R., and LIEBER, H. G. Non-Euclidean Geometry or Three Moons in Mathesis. 2d edition. Brooklyn, N. Y., H. G. L. R. Lieber, 1940. 40 pp.

MILES, E. R. C. See ELLIOTT, W. W.

PETERS, C. C., and VAN VOORHIS, W. R. Statistical Procedures and Their Mathematical Bases. New York and London, McGraw-Hill, 1940. 13+516 pp.

Tables of Circular and Hyperbolic Sines and Cosines. New York, Work Projects Administration, 1939. 405 pp.

TUKEY, J. W. Convergence and Uniformity in Topology. (Annals of Mathematics Studies, no. 2.) Princeton, University Press; London, Humphrey Milford and Oxford University Press, 1940. 9+90 pp.

VAN VOORHIS, W. R. See PETERS, C. C.

VAN DER WAERDEN, B. L. Moderne Algebra. Part 2. 2d revised edition. (Die Grundlehren der mathematischen Wissenschaften in Einzeldarstellungen, vol. 34.) Berlin, Springer, 1940. 8+224 pp.

WARREN, A. G. Mathematics Applied to Electrical Engineering. (Monographs on Electrical Engineering, vol. 9.) London, Chapman and Hall, 1939. 15+384 pp.

Wiley Trigonometric Tables, New York, Wiley; London, Chapman and Hall, 1940. 3+81 pp.